


BIOLOGY
Section I
53 Multiple-Choice Questions
5 Grid-In Questions
Time—90 Minutes

Directions: Each of the questions or incomplete statements below is followed by four suggested answers or completions. Select the one that is best in each case and then fill in the corresponding circle on the answer sheet.

1. A genetic counselor is consulted by a young man who is worried about developing Huntington's disease, an inherited disorder caused by a dominant allele of a single gene. The young man explains that his cousin was recently diagnosed with Huntington's disease, and the news has caused him to consider his own risk of developing the disorder. Which of the following questions will best help the genetic counselor to evaluate the risk of the young man developing Huntington's disease and transmitting it to his children?

(A) Were you and your cousin born in the same geographical area?
(B) Were your parents or grandparents ever diagnosed with Huntington's disease?
(C) Were you in physical contact with a person diagnosed with Huntington's disease?
(D) Were you ever exposed to substances that are suspected of being mutagens?




2. Over several years, bacteria were isolated from members of a human population and tested for antibiotic resistance. The percent of bacterial isolates that were found to be antibiotic resistant is presented in the graph above for each year of the study. Which of the following conclusions is best supported by the information presented in the graph?
- (A) The greatest increase in antibiotic resistance occurred from the first year to the third year of the study.
(B) Gene flow in the population occurred at the same rate for the entire duration of the eight-year study.
(C) Because evolution in bacterial populations occurs over millions of years, the results are due to random variation.
(D) Natural selection over multiple generations is favoring antibiotic resistance in the population of bacteria.

3. Lobe-finned fishes were present in the oceans of the world approximately 400 million years ago. The first tetrapods (vertebrates that had limbs and could move on land) date to about 365 million years ago. One hypothesis states that early tetrapods evolved from lobe-finned fishes. Which of the following is the best plan for testing the hypothesis?
- (A) Determining the number of fins found on lobe-finned fishes from different geographical locations
 - (B) Measuring the proportion of radioisotopes found in fossils of early tetrapods
 - (C) Comparing the arrangements of bones in the fins of lobe-finned fishes and limbs of the earliest tetrapods
 - (D) Sequencing DNA isolated from fossils of lobe-finned fishes to determine the size of the genome
4. Which of the following describes a metabolic consequence of a shortage of oxygen in muscle cells?
- (A) An increase in blood pH due to the accumulation of lactic acid
 - (B) No ATP production due to the absence of substrate-level phosphorylation
 - (C) A buildup of lactic acid in the muscle tissue due to fermentation
 - (D) A decrease in the oxidation of fatty acids due to a shortage of ATP

5. Prairie dogs are small mammals that live in large colonies in burrows in the ground. Prairie dogs that are near their own relatives when a predator approaches are much more likely to issue a warning bark than those that are near unrelated prairie dogs. The prairie dogs that hear a warning bark are more likely to hide in their burrows than to remain above ground. However, the prairie dog that gives the warning bark is putting itself at increased risk of being identified and killed by the predator.

Which of the following presents the most likely evolutionary explanation for the behaviors described?

- (A) The warning bark changes the behavior of the related prairie dogs nearby, allowing the prairie dog's family to have increased survival and reproductive success.
- (B) The barking prairie dog chooses to warn other prairie dogs, leading to more prairie dogs living above ground.
- (C) The barking prairie dog is alerting unrelated prairie dogs to the predator, so it is not giving any advantage to its own relatives.
- (D) The failure of the individual to bark when surrounded by unrelated prairie dogs ensures survival of the individual.



6. One model of a sustainable fisheries practice is for individual fish to be removed from a natural population at a rate equal to the highest possible growth rate of an ideal population. The graph above represents a population of bluefin tuna living along the Atlantic coast. At which labeled point in the graph is the population growth rate the highest?
- (A) I
 - (B) II
 - (C) III
 - (D) IV

Questions 7-11**EFFECT OF WATER-SOLUBLE POLLUTANTS ON MEMBRANE PERMEABILITY**





Treatment Group	Treatment Solution*	Mean Absorbance of 460 nm Light ($n = 5$)	2×Standard Error of the Mean (2×SEM)
I	70% isopropanol; 30% water	0.164	0.032
II	90% isopropanol; 10% water	0.125	0.100
III	50% acetone; 50% water	0.215	0.034
IV	70% acetone; 30% water	0.274	0.018
V	100% water	0.095	0.004
*Concentrations are reported as percent by volume.			

A student formulated a hypothesis that water-soluble pollutants damage living organisms by increasing the permeability of cellular membranes. To test the hypothesis, the student investigated the effect of isopropanol and acetone on beet root cells. The vacuoles of beet root cells contain large amounts of betacyanin, a water-soluble pigment that is released into the extracellular environment as a result of increased membrane permeability.


The student prepared identical samples of beet root tissue and incubated each sample for 15 minutes in the specific solution for that group. At the end of the incubation period, the student measured the absorbance of 460 nm light for each sample. A greater concentration of betacyanin in the solution surrounding the beet root cells results in a greater absorbance of 460 nm light. The results of the experiment are shown in the table above.

7. Which of the following is the dependent variable in the experiment?
- (A) The percent by volume of isopropanol in the treatment solutions
 - (B) The percent by volume of water in the treatment solutions
 - (C) The length of time each sample was incubated in the treatment solutions
 - (D) The absorbance of 460 nm light by the treatment solutions


8. Which of the following graphs is the most appropriate representation of the experimental data?



Questions 9-11 are on page 10.




9. The illustration above is a model of a typical beet root cell. Based on the experimental results, which of the following best represents the effect of acetone on the permeability of cellular membranes?




10. The student analyzed the data from the investigation and concluded that the estimate of the mean of one treatment group was unreliable. Which of the following identifies the treatment group most likely to have provided an unreliable estimate of the mean, and correctly explains why the estimate appears unreliable?
- (A) Treatment group II; it has a lower than expected mean absorbance and the largest standard error of the mean.
 - (B) Treatment group III; it has a higher than expected mean absorbance and the largest standard error of the mean.
 - (C) Treatment group IV; it has a higher than expected mean absorbance and the smallest standard error of the mean.
 - (D) Treatment group V; it has a lower than expected mean absorbance and the smallest standard error of the mean.
11. Based on the data from the investigation, which of the following is the best scientific question about organisms living in water that is polluted with organic solvents?
- (A) Do organisms without pigments have a selective advantage in polluted environments?
 - (B) Will organisms living in polluted environments exhibit detrimental effects from an increased permeability of their cellular membranes?
 - (C) Will organic solvents from the environment become incorporated into the cell membranes of organisms living in polluted environments?
 - (D) Should governments place more stringent restrictions on the release of organic solvents into aquatic ecosystems?

12. Within a forest ecosystem, there is a large amount of diversity among members of a warbler species. Of the following stages of meiosis illustrated for a typical cell, which contributes most to diversity among the warblers?


(A)




(B)



(C)



(D)



13. In an experiment, 100 mice were released into a field to which no other mice had access. Immediately after their release, a representative sample of mice was captured, their fur color was recorded, and they were returned to the field. After twenty years, a representative sample of mice was captured and the distribution of fur color was again recorded.

Fur Color	Initial Distribution (%)	Distribution after Twenty Years (%)
Brown	80	20
Gray	10	75
Black	10	5

Which of the following could best explain the change in fur color distribution, as shown in the table above?


- (A) The allele for black fur color is unstable, and over twenty years most of the black fur alleles mutated to become alleles for gray fur.
- (B) The field was primarily composed of light-colored soil and little vegetation, affording gray mice protection from predators.
- (C) Sexual selection led to increased mating frequency of black and brown versus gray and brown.
- (D) The gray mice were harder to catch, and so were underrepresented in the twenty-year sample.

14. Liver cells manufacture glycoproteins, while adipose cells store fat. Which of the following subcellular structures is likely to be more prominent in liver cells than in adipose cells?

- (A) Nucleus
- (B) Golgi apparatus
- (C) Cytoskeleton
- (D) Plasma membrane

15. Intravenous (IV) solutions administered to patients are normally isotonic. Which of the following is most likely if an IV of distilled water is administered to a patient?

- (A) The cells that are exposed to hypotonic solutions will shrink as a result of salt moving into the blood.
- (B) The liver will secrete additional bile salts into the blood to raise the tonicity of the administered fluid.
- (C) The cells that are exposed to hypotonic solutions will expand as water moves osmotically into the cells from the blood.
- (D) The patient's respiration rate will slow to compensate for the higher levels of circulating blood.




16. The diagram above represents a typical rod-shaped bacterium. Which of the following best describes a feature shown in the diagram that is unique to archaea and bacteria?

- (A) The organism is surrounded by a cell wall.
- (B) The organism contains ribosomes.
- (C) The organism does not have a nuclear membrane surrounding its genetic material.
- (D) The organism is not capable of making or providing itself with ATP.

17. Which of the following best supports the claim that organisms of different domains share a common ancestor?

- (A) Photosynthesis occurs in chloroplasts, and the citric acid cycle occurs in mitochondria.
- (B) Glycolysis occurs in both prokaryotic cells and eukaryotic cells.
- (C) Introns are present in eukaryotic DNA but not in prokaryotic DNA.
- (D) Errors in DNA synthesis provide some of the genetic variation in a population.

18. The graph below shows a model of population changes over time in a predator and its prey.



Which of the following data would be most helpful in predicting possible disruptions to the relationship between the predator and its prey?

- (A) Time of day, because as days become shorter, nighttime hunting increases
- (B) Average birth weight of predators, because heavier offspring will begin hunting at younger ages
- (C) Numbers of additional predator species, because additional predators will increase competition for prey
- (D) Prey metabolic rates, because predators are less successful against prey with higher metabolic rates

Questions 19-22

The three-spined stickleback (*Gasterosteus aculeatus*) is a small fish found in both marine and freshwater environments. Marine stickleback populations consist mostly of individuals with pronounced pelvic spines, as shown in Figure 1. Individuals in freshwater stickleback populations, on the other hand, typically have reduced pelvic spines, as shown in Figure 2.




Figure 1. Marine stickleback





Figure 2. Freshwater stickleback

As represented in Figure 1 and Figure 2, the phenotypic difference between marine and freshwater sticklebacks involves *Pitx1*, a gene that influences the formation of the jaw, pituitary gland, and pelvic spine. Enhancer sequences upstream of the *Pitx1* genetic locus regulate expression of the *Pitx1* gene at the appropriate times and in the appropriate tissues during development. Previous studies have found that a mutation in the hindlimb enhancer interferes with the formation of a pronounced pelvic spine.


19. Which of the following best describes how sticklebacks in the same population with identical copies of the *Pitx1* gene can still show phenotypic variation in the pelvic spine character?
- (A) The *Pitx1* gene is carried on different chromosomes in different individuals.
 - (B) Expression of the *Pitx1* gene is affected by mutations at other genetic loci.
 - (C) The genetic code of the *Pitx1* gene is translated differently in males and females.
 - (D) The subcellular location of the *Pitx1* gene changes when individuals move to a new environment.
20. A mutation that affects *Pitx1* gene function in all tissue types is most likely to be at which of the following genetic loci?
- (A) Hindlimb enhancer
 - (B) Pituitary enhancer
 - (C) Jaw enhancer
 - (D) Promoter
21. In sticklebacks, which of the following is most likely to occur if the jaw enhancer is disabled instead of the hindlimb enhancer?
- (A) The jaw and a pronounced pelvic spine develop normally because the *Pitx1* gene is expressed in both developing tissues.
 - (B) Neither the jaw nor a pronounced pelvic spine develop normally because there is no *Pitx1* gene expression in either developing tissue.
 - (C) The jaw develops normally, but a pronounced pelvic spine does not develop because the *Pitx1* gene is expressed in the developing jaw but not in the developing pelvis.
 - (D) The jaw does not develop normally, but a pronounced pelvic spine does develop because the *Pitx1* gene is expressed in the developing pelvis but not in the developing jaw.
22. Which of the following describes a possible selective mechanism to explain why freshwater sticklebacks typically have reduced pelvic spines?
- (A) Reduced pelvic spines increase the likelihood of escaping predators in freshwater environments.
 - (B) Reduced pelvic spines expand the range of foods that can be used as sources of energy in freshwater environments.
 - (C) Reduced pelvic spines result in males becoming effectively invisible to potential mating partners in freshwater environments.
 - (D) Reduced pelvic spines allow individual fish to pass through narrow waterways on their way to marine environments.

23. *Hydrangea macrophylla* is a species of plant with blue or pink flowers. Flower color in *Hydrangea macrophylla* plants is thought to be determined primarily by soil conditions rather than by inherited differences. Which of the following will provide the most direct evidence that flower color in *Hydrangea macrophylla* is due primarily to soil conditions?

- (A) Comparing DNA sequence information for *Hydrangea macrophylla* to DNA sequence information for a different species of flowering plant
- (B) Performing a genetic cross between a pink-flowered plant and a blue-flowered plant and then determining the ratio of phenotypes in the offspring
- (C) Growing cuttings from the same *Hydrangea macrophylla* plant under controlled conditions that vary only with regard to soil pH
- (D) Analyzing fossilized remains of an ancestral species that grew in geographic areas similar to the habitats of modern *Hydrangea macrophylla*



24. Based on morphological characteristics, the phylogenetic tree above has been developed for species I, II, III, and IV. DNA sequencing has recently been completed for a particular gene found in all four species. The sequencing will provide additional information for answering which of the following questions?
- (A) Is species I the common ancestor of species II and III?
 - (B) Does species I contain the greatest number of vestigial structures?
 - (C) Are species III and IV the most closely related?
 - (D) Should the positions of species III and IV be reversed?
25. The transmission of information from sensory neurons to interneurons typically involves the
- (A) exchange of electrons between specialized proteins embedded in the plasma membrane of the sensory neuron to the plasma membrane of the interneuron
 - (B) release of chemical messengers into the space between the axon of the sensory neuron and the plasma membrane of the interneuron
 - (C) transfer of a phosphate group from ATP in the sensory neuron to a protein substrate in the interneuron
 - (D) flow of protons down an electrochemical gradient through a gated channel between the sensory neuron and the interneuron



26. The brain coordinates the circulatory and respiratory systems of the human body. The control of breathing, for example, involves neural pathways among the structures represented in the figure above. One important stimulus in the control of breathing is an increase in blood CO_2 concentration, which is detected as a decrease in blood pH. Which of the following best describes the physiological response to an overall increase in cellular respiration in the body?
- (A) In response to depleted blood CO_2 levels, the pH sensors send signals directly to the rib muscles, resulting in an increase in the rate of CO_2 uptake by the lungs and a decrease in CO_2 utilization by the brain.
 - (B) In response to low blood pH, the pH sensors send a signal to the brain, which then sends a signal to the diaphragm, resulting in an increased rate of breathing to help eliminate excess blood CO_2 .
 - (C) In response to high blood pH, the pH sensors send a signal directly to the lungs, resulting in a slower rate of breathing, and the lungs send a signal back to the heart once CO_2 availability has been restored.
 - (D) In response to an increased rate of breathing, the rib muscles send a signal to the brain, which then sends a signal to the heart, resulting in a decrease in heart activity and slower flow of blood through the body.

27. Thyroxin is a hormone that increases metabolic activities within various tissue targets. Low levels of circulating thyroxin trigger the secretion of thyroid-stimulating hormone (TSH) from the anterior pituitary. TSH secretion then stimulates thyroxin production and release by the thyroid gland. The increased level of circulating thyroxin inhibits further secretion of TSH from the anterior pituitary.

Based on the information provided, which of the following can most likely be concluded about the TSH–thyroxin loop?

- (A) A person taking thyroxin to supplement low thyroxin secretion will produce more TSH.
- (B) Increased thyroxin production would cause elevated ribosomal activity in the anterior pituitary.
- (C) The structure of the loop would lead to elevated thyroid and tissue activity due to positive feedback.
- (D) The feedback mechanism would maintain relatively constant levels of thyroxin throughout tissue targets.

28. The Stanley Miller apparatus demonstrated that organic molecules could assemble spontaneously in an environment lacking free oxygen and containing water, methane, and ammonia in the presence of an abundant energy source, such as an electric discharge. The research was considered supportive of the organic soup hypothesis, which states that the primitive atmosphere provided inorganic precursors from which organic molecules could have been synthesized in the presence of an energy source. Based on subsequent research, the primordial atmosphere was determined to contain less methane and more carbon dioxide. The new data about the composition of the early atmosphere had which of the following effects on origin-of-life hypotheses?

- (A) Miller's work was discarded as not relevant to an origin-of-life hypothesis.
- (B) Miller's work was used to refute the new hypothesis about the composition of the atmosphere.
- (C) The organic soup hypothesis was abandoned as no longer being supportable.
- (D) A new organic soup hypothesis was proposed to account for the new data about the atmosphere.

Questions 29-33

Ascorbic acid (vitamin C) is an organic molecule necessary for the health of plants and animals. The majority of animals, including most mammals, synthesize ascorbic acid from organic precursors, but some primates are unable to synthesize ascorbic acid and must instead acquire it from dietary sources, such as certain fruits and vegetables.

The *L-gulonolactone oxidase (GULO)* gene encodes an enzyme that catalyzes a required step in the biosynthesis of ascorbic acid. Most mammals carry a functional copy of the *GULO* gene, but some primates carry only a *GULO* pseudogene, which is a nonfunctional variant.

A comparison of *GULO* genes and *GULO* pseudogenes from different animals can provide insight into the evolutionary relatedness of the animals. In Table I, selected members of some mammalian groups are listed, along with an indication of their ability to synthesize ascorbic acid. Table II shows an alignment of amino acid coding sequences from homologous regions of the *GULO* genes and *GULO* pseudogenes of the organisms listed in Table I. Figure 1 represents the universal genetic code.

TABLE I: SELECTED MAMMALIAN GROUPS

Group	Selected Members	Biosynthesis of Ascorbic Acid
Nonprimate mammals	Elephant, mouse	Yes
Primate mammals	Lemur	Yes
	Orangutan, chimpanzee	No
	Human	No

TABLE II: DNA SEQUENCE ALIGNMENT*

	Relative Positions of Nucleotides in Nontemplate (Coding) Sequence																										
	1 (5')	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27 (3')
Elephant	G	A	C	A	C	C	C	A	T	C	T	G	A	A	G	A	A	G	T	C	G	G	A	A	T	A	C
Mouse	G	A	C	A	G	C	C	A	C	C	T	G	A	A	G	A	A	G	T	C	T	G	A	G	T	A	C
Lemur	G	A	C	A	G	C	C	A	C	C	T	G	A	A	G	A	G	G	T	C	C	G	A	G	T	A	C
Orangutan	G	A	C	A	G	C	-	A	T	T	G	G	A	A	G	A	A	A	T	C	T	G	A	G	G	A	C
Chimp	G	A	C	A	G	C	-	A	T	T	G	G	A	A	G	A	A	A	T	C	T	G	A	G	G	A	C
Human	G	A	C	A	G	C	-	A	T	T	G	G	A	A	G	A	A	A	T	C	T	G	A	G	G	A	C

*For each DNA segment, the alternating shaded and unshaded nucleotides indicate the triplet codons of the open reading frame, shown from left (5') to right (3') as the nontemplate (coding) strand. An “-” indicates the absence of a nucleotide.

		Second Base				
		U	C	A	G	
5' First Base	U	UUU } Phe	UCU } Ser	UAU } Tyr	UGU } Cys	U
		UUC }	UCC }	UAC }	UGC }	C
		UUA } Leu	UCA }	UAA stop	UGA stop	A
		UUG }	UCG }	UAG stop	UGG Trp	G
	C	CUU } Leu	CCU } Pro	CAU } His	CGU } Arg	U
		CUC }	CCC }	CAC }	CGC }	C
		CUA }	CCA }	CAA } Gln	CGA }	A
		CUG }	CCG }	CAG }	CGG }	G
	A	AUU } Ile	ACU } Thr	AAU } Asn	AGU } Ser	U
		AUC }	ACC }	AAC }	AGC }	C
		AUA }	ACA }	AAA } Lys	AGA } Arg	A
		AUG Met	ACG }	AAG }	AGG }	G
	G	GUU } Val	GCU } Ala	GAU } Asp	GGU } Gly	U
		GUC }	GCC }	GAC }	GGC }	C
		GUA }	GCA }	GAA } Glu	GGA }	A
		GUG }	GCG }	GAG }	GGG }	G

Figure 1. Universal genetic code.

Questions 29-33 are on pages 22 and 23.





Item 29 was not scored.


30. Comparison of DNA sequences in Table II suggests that a functional *GULO* gene in lemurs can have a G, C, or T at position 21 but only a G at position 22. Which of the following pairs of predictions is most helpful in explaining the discrepancy?

	A substitution at position 21 would result in	A substitution at position 22 would result in
(A)	No change to the protein	A premature stop codon or an amino acid with different biochemical characteristics
(B)	A different amino acid	A premature stop codon or an amino acid with different biochemical characteristics
(C)	No change to the protein	A frame shift producing an inactive protein
(D)	An amino acid with different biochemical characteristics	No transcription of the gene

31. Which of the following is the most likely effect of the mutation at nucleotide position 7 in the *GULO* gene of humans?
- (A) The mutation results in the deletion of the *GULO* gene, so no polypeptide can be translated.
 - (B) The deletion of the single nucleotide causes a frame shift, changing the primary structure downstream of the mutation and resulting in a nonfunctional protein.
 - (C) The point mutation causes a substitution of the amino acid isoleucine (Ile) for histidine (His) at position 7, resulting in a protein with higher than normal activity.
 - (D) The substitution of a single nucleotide in the *GULO* coding region results in a stop codon. This results in a smaller nonfunctional protein.
32. Lemurs are primates that live on the island of Madagascar off the coast of Africa. Lemurs have a functional *GULO* gene and are able to produce ascorbic acid. However, primates that live in other places (e.g., humans, chimpanzees, and orangutans) have a *GULO* pseudogene and are unable to produce ascorbic acid. Which of the following best explains the genetic variation among primate species?
- (A) Food sources on the island of Madagascar were deficient in ascorbic acid, which caused the primates to migrate to the mainland where ascorbic acid-rich foods are more widely available.
 - (B) Human activity on Madagascar, including poaching, drove the other primates into extinction on the island.
 - (C) Food sources where non-Madagascar primates lived provided ample ascorbic acid in the diet, which removed the selective pressure for maintaining a functional *GULO* gene.
 - (D) Food sources on the island of Madagascar were deficient in ascorbic acid, which caused lemurs to adapt by developing the ability to produce an active *GULO* protein.

33. Which of the following phylogenetic trees best illustrates (with the X) the point at which the mutation in the *GULO* gene most likely occurred during the evolutionary history of these organisms?





34. The cells lining the respiratory tract of pigs have receptors for both avian and human influenza viruses. Based on the model above, which of the following best describes the origin of the new strain of human influenza virus?
- (A) The new viral strain inherited a mixture of genetic material from both avian influenza virus and human influenza virus.
 - (B) The new viral strain inherited RNA molecules from the avian influenza virus and packaged them inside the human influenza virus membrane.
 - (C) The new viral strain inherited a mutant DNA molecule from the pig respiratory tract cell.
 - (D) The new viral strain inherited an RNA molecule that had recombined with a DNA molecule from the pig respiratory tract cell.

35. In the “RNA world” model for the origin of life, the first protocells (probiotics) relied on RNA, not DNA, for information storage and transmission. Which of the following could best be considered evidence in support of the RNA world model?

- (A) DNA is more stable than is RNA.
- (B) Some RNA molecules in contemporary cells have catalytic function.
- (C) RNA is so complex in both structure and function that it must have preceded DNA.
- (D) RNA has secondary structure whereas DNA does not.

36. Which of the following pieces of evidence best supports the hypothesis that birds and crocodilians (crocodiles and alligators) are more closely related to each other than they are to other organisms?

- (A) Modern birds and crocodilians utilize different niches within the same geographical region.
- (B) Like turtles and snakes, modern birds and crocodilians build nests in which they lay shelled eggs.
- (C) The skulls of birds and crocodilians have an opening that is absent in the skulls of other reptiles.
- (D) Fossil evidence indicates that both modern birds and crocodilians originated during the Jurassic Period.

Cellular Shape	Ratio of Surface Area to Volume
Cube	1.7
Regular Tetrahedron	0.5
Cylinder	0.9
Sphere	0.6

37. In an experiment, the efficiency of oxygen exchange across the plasma membrane is being assessed in four artificial red blood cells. The table above lists some properties of those artificial cells. Other conditions being equal, which artificial cell is predicted to be the most efficient in exchanging oxygen with the environment by diffusion?

- (A) The cuboidal cell
- (B) The tetrahedral cell
- (C) The cylindrical cell
- (D) The spherical cell

38. To test the hypothesis that a particular plant synthesizes storage lipids by using glyceraldehyde 3-phosphate (G3P) from photosynthesis, a researcher plans to use radiolabeled precursors to track the molecules through the biosynthetic pathway. Which of the following radiolabeled precursors is most appropriate for the researcher to use?
- (A) ^{15}N -labeled N_2 , because atmospheric nitrogen is fixed to amino acids by photosynthesis
 - (B) ^{14}C -labeled CO_2 , because atmospheric carbon is fixed to carbohydrates by photosynthesis
 - (C) ^{35}S -labeled methionine, because amino acids are incorporated into lipids during photosynthesis
 - (D) ^{32}P -labeled phosphate, because lipids are stored in plants as phospholipids

- Diagram illustrating a point mutation in DNA:
- Top strand (Wild-type): ...CAGACTAGGGCCAGAGCCTTTAGCTTGTCCGTAAAAGAGC...
- Bottom strand (Wild-type): ...GTCCTGATCCCGGCTCTGGGAAATTCGAAACAGGACGATTTTCTCTGCTG...
- Mutation event: DNA Damage, Imperfect Repair
- Bottom strand (Mutant): ...CAGACTAGGGCCAGAGCCTTTA**A**CTTGTCCGTAAAAGAGC...
- The diagram shows a point mutation where the 10th nucleotide of the bottom strand (T) is replaced by an A (Adenine), resulting in a C to A transition. This mutation is labeled as "DNA Damage, Imperfect Repair".

(A) ... CAGACTAGGGCCAGAGCCTTTA**G**CTTGTCCGTAAAAGAGC ... Normal
 ... G L C L G A L C C C G G L C L C G G A A A L C G A A C A G G C A L L L L C L C G ...
 and
 ... CAGACTAGGGCCAGAGCCTTTA**G**CTTGTCCGTAAAAGAGC ... Normal
 ... G L C L G A L C C C G G L C L C G G A A A L C G A A C A G G C A L L L L C L C G ...

(B) ... CAGACTAGGGCCAGAGCCTTTA**A**CTTGTCCGTAAAAGAGC ... Mutant
 ... G L C L G A L C C C G G L C L C G G A A A L L G A A C A G G C A L L L L C L C G ...
 and
 ... CAGACTAGGGCCAGAGCCTTTA**G**CTTGTCCGTAAAAGAGC ... Normal
 ... G L C L G A L C C C G G L C L C G G A A A L C G A A C A G G C A L L L L C L C G ...

(C) ... CAGACTAGGGCCAGAGCCTTTA**A**CTTGTCCGTAAAAGAGC ... Mutant
 ... G L C L G A L C C C G G L C L C G G A A A L L G A A C A G G C A L L L L C L C G ...
 and
 ... CAGACTAGGGCCAGAGCCTTTA**A**CTTGTCCGTAAAAGAGC ... Mutant
 ... G L C L G A L C C C G G L C L C G G A A A L L G A A C A G G C A L L L L C L C G ...

(D) ... CAGACTAGGGCCAGAGCCTTTA**A**CTTGTCCGTAAAAGAGC ... Mutant
 ... G L C L G A L C C C G G L C L C G G A A A L C G A A C A G G C A L L L L C L C G ...
 and
 ... CAGACTAGGGCCAGAGCCTTTA**A**CTTGTCCGTAAAAGAGC ... Mutant
 ... G L C L G A L C C C G G L C L C G G A A A L C G A A C A G G C A L L L L C L C G ...

Questions 40-44

Iron is an essential nutrient that is acquired by organisms from the environment. When intracellular levels of iron are relatively high, living cells synthesize an iron-storage protein called ferritin.

The induction of ferritin synthesis by iron was investigated in rats. Figure 1 shows the results of an experiment in which cellular levels of ferritin protein were measured in the presence or absence of iron and actinomycin D, a drug that inhibits transcription.

Figure 2 shows the results of an experiment in which cellular levels of ferritin protein were measured in the presence or absence of iron and cycloheximide, a drug that inhibits translation.

Iron	–	+	–	+
Actinomycin D	–	–	+	+




Figure 1. Ferritin Protein Levels

Iron	–	+	–	+
Cycloheximide	–	–	+	+





Figure 2. Ferritin Protein Levels

40. Which of the following biotechnology approaches could be used to identify ferritin mRNA in a sample of total cellular RNA?
- (A) RNA samples could be directly cloned into a DNA plasmid, grown in bacteria, and tested for the ability to bind iron.
 - (B) RNA samples could be separated by size using agarose gel electrophoresis and incubated with labeled single-stranded DNA molecules that are complementary to the ferritin mRNA.
 - (C) RNA samples could be converted to protein and subsequently cut with a restriction endonuclease that recognizes DNA sequences.
 - (D) RNA samples could be examined under a high-power microscope to visually identify the ferritin mRNA.
41. In individuals with iron-rich diets, intracellular levels of iron are higher than in individuals with iron-poor diets. Which of the following best explains how organisms regulate ferritin protein levels in response to available iron?
- (A) Translation of ferritin mRNA is inhibited in the presence of iron.
 - (B) Translation of ferritin mRNA is stimulated by iron.
 - (C) Ferritin protein is more stable in the presence of iron.
 - (D) Ferritin protein is less stable in the presence of iron.
42. The data can best be used to support which of the following claims about the mechanism for regulating ferritin gene expression?
- (A) Iron increases ribosome binding to ferritin mRNA.
 - (B) Iron causes the ferritin mRNA to be degraded.
 - (C) Iron increases the effect of actinomycin D on RNA polymerase.
 - (D) Iron decreases the ability of ferritin mRNA to bind to ribosomes.
43. The gene sequences responsible for the iron-mediated changes in ferritin protein levels are highly conserved and are called iron response elements (IREs). IREs have been observed in a number of genes involved in iron transport and metabolism. The IRE sequences found in the ferritin gene are found in all multicellular organisms, whereas other nonferritin IRE sequences are found only in certain vertebrate organisms. Which of the following is the best explanation for the observations?
- (A) Ferritin IREs arose in the common ancestor of multicellular organisms, whereas nonferritin IREs arose later in specific evolutionary lineages.
 - (B) Nonferritin IREs arose at different times in the common ancestor of multicellular organisms.
 - (C) Ferritin IREs arose in diverse organisms by convergent evolution, while the nonferritin IREs arose by recombination events between closely related species.
 - (D) Ferritin IREs have evolved independently in different evolutionary lineages, whereas nonferritin IREs arose in the common ancestor of multicellular organisms.
44. Hereditary hemochromatosis (HHC) is an iron storage disease that results in reduced uptake of iron into cells. Based on the data, which of the following is the most likely effect of HHC on ferritin protein levels?
- (A) Elevated levels of ferritin protein because there is more iron in the blood of affected individuals
 - (B) Reduced levels of ferritin protein because there is less iron in the cytosol of affected individuals
 - (C) Elevated levels of ferritin protein because ferritin replaces the mutant protein in affected individuals
 - (D) Reduced levels of ferritin protein because iron stimulates ferritin breakdown



45. The figure above shows a series of microscope images taken over time of ovalbumin production in chick oviduct cells in response to stimulation with estrogen. The ovalbumin protein was detected using a fluorescent dye as shown by the white areas in the time-lapse sequence.

The microscope images indicate that ovalbumin

- (A) is synthesized on soluble ribosomes and packaged into lysosomes, from which it is secreted into the oviduct
- (B) is synthesized on ribosomes bound to the rough endoplasmic reticulum, and then secreted directly into the oviduct
- (C) crosses the endoplasmic reticulum membrane as it is translated, then moves to the Golgi apparatus, then to vesicles from which it is secreted
- (D) crosses the Golgi apparatus membrane post-translationally, is taken up by the Golgi apparatus and endoplasmic reticulum, then is packaged into secretory vesicles

46. Freshwater bony fish secrete large volumes of very dilute urine. Many marine bony fish secrete small volumes of concentrated urine. Which of the following best explains the differences in nitrogenous-waste excretion between freshwater and marine bony fish?

- (A) The differences arose during divergence, as fish from a common ancestor faced different selective pressures in environments of different osmolarity.
- (B) The differences resulted from the evolution of kidneys in independent evolutionary lineages that were adapted to different environmental conditions.
- (C) The differences are accidents of evolutionary divergence and are unrelated to the osmotic differences in freshwater and marine environments.
- (D) The differences arise in each generation because of the physiological adjustment of individual fish to fluctuating salt concentrations in the environments in which they live.

47. Malaria is caused by several different species of *Plasmodium*, a protozoan parasite. *Plasmodium* resistance to the common drugs used to treat malaria has increased in recent years.

In a scientific study, *Plasmodium* samples were analyzed in blood drawn from a large number of infected patients before drug treatment and subsequently from the subset of infected patients with drug-resistant *Plasmodium*. DNA sequences of four different *Plasmodium* genes thought to be involved in resistance were compared between samples from patients with drug-sensitive *Plasmodium* and patients with drug-resistant *Plasmodium*.

Which of the following best supports the hypothesis that preexisting mutations confer drug resistance?

- (A) Drug resistance can sometimes be reversed by simultaneous treatment with a drug that inhibits the drug-pumping mechanism of *Plasmodium*.
- (B) Some of the mutations that occur most frequently in samples from patients treated with antimalarial drugs are detectable in *Plasmodium* samples collected prior to use of antimalarial drugs.
- (C) Continued use of the same antimalarial drugs leads to increased *Plasmodium* resistance.
- (D) The same mechanism of drug resistance in *Plasmodia* isolated from separate populations of patients resulted from an accumulation of new genetic mutations.

Questions 48-50

Insulin, a hormone secreted by pancreatic cells, stimulates glucose uptake in skeletal muscle cells by mobilizing glucose transporter proteins (GLUT4) to the plasma membrane. As depicted in Figure 1, binding of insulin to the insulin receptor triggers an intracellular signaling cascade in which certain molecules activate other molecules in a relay of the hormone signal to cell targets. One outcome of the signaling cascade is mobilization of GLUT4 from vesicle storage sites in the cytoplasm to sites at the cell surface, where GLUT4 allows glucose to enter the cell.




Figure 1. Insulin signaling in muscle cells

In type 2 diabetes, the cellular response to insulin is disrupted, and individuals with type 2 diabetes cannot properly regulate their blood glucose levels. In an investigation of the insulin signaling pathway, samples of skeletal muscle were isolated from individuals who have type 2 diabetes and from individuals who do not. The results of several experiments that were performed on the muscle samples are shown in Figure 2, Figure 3, and Figure 4.




Figure 2. Insulin-stimulated glucose uptake




Figure 3. Insulin receptor activation




Figure 4. IRS-1 activation

48. Which of the following is a valid interpretation of the experimental results that explains how individuals with type 2 diabetes differ from individuals without diabetes?
- (A) The relatively low levels of glucose uptake in individuals with type 2 diabetes indicate that mobilization of GLUT4 to the cell surface is reduced in muscle cells of those individuals.
 - (B) The relatively low levels of glucose uptake in individuals with type 2 diabetes indicate that no functional GLUT4 protein is produced in the muscle cells of those individuals.
 - (C) The absence of activated insulin receptors in individuals with type 2 diabetes indicates that no insulin is secreted by the pancreatic cells of those individuals.
 - (D) The absence of activated IRS-1 in individuals with type 2 diabetes indicates that no functional insulin receptor protein is produced in the muscle cells of those individuals.
49. Based on the experimental results, which of the following describes the most likely defect in muscle cells of patients with type 2 diabetes?
- (A) Insulin receptor proteins do not reach the cell surface.
 - (B) Insulin does not activate its receptor.
 - (C) IRS-1 activation is reduced at high insulin concentrations.
 - (D) GLUT4 blocks glucose from entering cells.
50. Based on the information presented, which of the following genetic changes in an individual without diabetes is most likely to result in a disrupted cellular response to insulin signaling similar to that of an individual with type 2 diabetes?
- (A) A deletion in the gene encoding the insulin receptor that removes only the cytoplasmic domain of the protein
 - (B) Duplication of the gene encoding a PI-3 kinase that results in synthesis of a muscle-specific variant of the enzyme in skin cells as well as in muscle cells
 - (C) A mutation in the gene encoding IRS-1 that causes the protein to be active in muscle cells even in the absence of insulin signaling
 - (D) Insertion of a small segment of DNA into the promoter of the *Glut4* gene that results in increased synthesis of GLUT4 proteins in muscle cells

51. Trypsinogen is split by the enzyme enterokinase to form an activated molecule of the protease trypsin. Which of the following would confirm that the activation of trypsin is an example of how a positive feedback mechanism can amplify a biological process?
- (A) The activated trypsin enzyme can use enterokinase as a substrate.
 - (B) The trypsin produced by the reaction is capable of splitting and activating additional trypsinogen molecules.
 - (C) If levels of trypsin were to get too high, the trypsin molecules would inhibit the enzyme enterokinase.
 - (D) Each mRNA molecule that codes for trypsinogen can be translated repeatedly to form many peptide molecules.
52. Cancer can result from a variety of different mutational events. Which of the following is LEAST likely to result in the initiation of a cancerous tumor?
- (A) A receptor mutation results in activation of a cell-division pathway in the absence of the appropriate ligand.
 - (B) A mutation results in the loss of the ability to produce a tumor-suppressor protein.
 - (C) A defect in a cell-cycle checkpoint prevents a cell from entering the S phase.
 - (D) At the anaphase checkpoint, separation of chromatids occurs without all centromeres being attached to kinetochore microtubules from both poles.

53. Which of the following statements is true regarding the movement of substances across cell membranes?
- (A) Ions are unable to move through the phospholipid bilayer because the nonpolar tail regions of the phospholipids are hydrophobic.
 - (B) Ions are able to move through the phospholipid bilayer because the polar head regions of the phospholipids are charged.
 - (C) Water is able to move through the phospholipid bilayer because the nonpolar tail regions of the phospholipids are charged.
 - (D) Water is unable to move through the phospholipid bilayer because the polar head regions of the phospholipids are charged.

Determine the correct answer for each question and enter it in the grid on page 3 of the answer sheet. Use the following guidelines for entering your answers.

- Start your answer in any column, space permitting. Unused columns should be left blank.
- Write your answer in the boxes at the top of the grid and fill in the corresponding circles. Mark only one circle in any column. You will receive credit only if the circles are filled in completely.
- Provide your answer in the format specified by the question. The requested answer may be an integer, a decimal, or a fraction, and it may have a negative value.
- To enter a fraction, use one of the division slashes to separate the numerator from the denominator, as shown in the example below. Fractions only need to be reduced enough to fit in the grid.
- Do not enter a mixed number, as this will be scored as a fraction. For example, 2 1/2 (two and one-half) will be scored as 21/2 (twenty-one halves).

Fraction answer: $-2/10$
(does not have
to be reduced)

	5	0	2	4	
−	.	.	.	.	.
0					
1					
2					
3					
4					
5					
6					
7					
8					
9					

		5	0	2	4
−	.	/	/	/	.
0	0	●	0	0	
1	1	1	1	1	
2	2	2	●	2	
3	3	3	3	3	
4	4	4	4	●	
5	●	5	5	5	
6	6	6	6	6	
7	7	7	7	7	
8	8	8	8	8	
9	9	9	9	9	


-		4	.	1	3
●	○	/ ○	/ ●	/ ○	○
	(0)	(0)	(0)	(0)	(0)
	(1)	(1)	(1)	●	(1)
	(2)	(2)	(2)	(2)	(2)
	(3)	(3)	(3)	(3)	●
	(4)	●	(4)	(4)	(4)
	(5)	(5)	(5)	(5)	(5)
	(6)	(6)	(6)	(6)	(6)
	(7)	(7)	(7)	(7)	(7)
	(8)	(8)	(8)	(8)	(8)
	(9)	(9)	(9)	(9)	(9)

[illegible]

121. The internal temperature of different organisms was determined under various conditions, as shown in the table below.


Organism	Thermoregulation Strategy	Measurement Conditions	Internal Temperature
Butterfly	Ectotherm	Resting in the shade	15°C
Butterfly	Ectotherm	Flying in the sunlight	23°C
Horse	Endotherm	Resting in the shade	42°C
Horse	Endotherm	Galloping in a field	43°C
Goose	Endotherm	Sitting on a nest	41°C
Goose	Endotherm	Flying above treetops	??°C

Predict the internal temperature (in °C) of a goose that is flying just above the treetops when the external temperature is 7°C. Give your answer to the nearest whole number.



122. The graph above shows the genotype frequencies for a particular genetic locus at different allele frequencies in a population. Using the graph, determine the frequency of the dominant allele in the population if the frequency of homozygous recessive individuals in the population is 0.75. Enter your answer as a value between zero and one to the nearest hundredth.

123. As part of a study investigating the temperature profile of a temperate-zone lake, water temperature measurements were made at different water depths at the same time on the same day. The results are presented in the graph below.



According to the data, a strict ectotherm that acclimates at 5 meters and again at 15 meters will be exposed to how many degrees change in temperature? Provide your answer to the nearest tenth of a degree.

124. Phenylketonuria (PKU) is an inherited disease caused by an autosomal recessive allele.

If two individuals who are carriers of PKU have two children, what is the probability that neither child will have PKU? Give your answer as a fraction or decimal.

125. The effect of temperature on the respiration rate of heterotrophic bacteria in a shallow, slow-moving freshwater stream was investigated by performing the following experiment. The water temperature of the stream was monitored over a continuous 24-hour period (starting at time = 0 hours). The respiration rates of bacteria living in the streambed were determined at 6 hours and at 9 hours. The results are presented below.




Figure 1. Water Temperature




Figure 2. Respiration Rate

Based on the information provided, what was the change (in °C) in water temperature associated with the change in respiration rate shown in Figure 2 ? Give your answer to one decimal place.

BIOLOGY

Section II

8 Questions

Planning Time—10 minutes

Writing Time—80 minutes

Directions: Questions 1 and 2 are long free-response questions that require about 22 minutes each to answer and are worth 10 points each. Questions 3–8 are short free-response questions that require about 6 minutes each to answer. Questions 3–5 are worth 4 points each and questions 6–8 are worth 3 points each.

Read each question carefully and completely. Write your response in the space provided for each question. Only material written in the space provided will be scored. Answers must be written out in paragraph form. Outlines, bulleted lists, or diagrams alone are not acceptable.

Question 1 is on the following page.

1. To determine the likelihood that a patient with a bacterial infection who is in the intensive care unit (ICU) of a hospital could be successfully treated with antibiotic therapies, researchers investigated the frequency of antibiotic resistance in bacteria isolated from patients in ICUs in the United States from 1994 to 2000. The data from the study are shown in the table, along with the overall change in the frequency of bacterial resistance in patient samples from 1994 to 2000.

INCIDENCE OF ANTIBIOTIC RESISTANCE IN PATIENTS ADMITTED TO ICUs FROM 1994 TO 2000

Antibiotic	Percent of All Samples Demonstrating Antibiotic Resistance in 2000	Overall Change in Percent Resistance to Antibiotics from 1994 to 2000
Ciprofloxacin	19	+10
Gentamicin	22	+6
Ampicillin	65	+3
Imipenem	11	+1
Tazobactam	22	0

- (a) A patient was admitted to the ICU in December 2000 with an uncharacterized bacterial infection. Based on the data from the study, **identify** the following.
- The antibiotic that was most likely to be effective for treating the infection
 - The antibiotic that was least likely to be effective for treating the infection
- (b) Using the data, **provide justification** for the claim that antibiotic resistance may arise in bacterial species.
- (c) **Describe** THREE different processes by which antibiotic resistance develops in individual bacterial cells.
- (d) **Describe** TWO features of natural selection that lead to the spread of antibiotic resistance in a population over many generations. **Give ONE reason** that the development of antibiotic resistance is an example of evolution.
- (e) Giving low doses of antibiotics to beef cattle to control the population sizes of intestinal bacteria can result in more efficient uptake of food materials by the cattle. **Pose** an ethical question that the Food and Drug Administration (FDA) should consider when deciding whether to approve this type of antibiotic use in beef cattle.

THIS PAGE MAY BE USED FOR TAKING NOTES AND PLANNING YOUR ANSWERS.

NOTES WRITTEN ON THIS PAGE WILL NOT BE SCORED.

WRITE ALL YOUR RESPONSES ON THE LINED PAGES.

GO ON TO THE NEXT PAGE.


[illegible]

[illegible]

[illegible]

[illegible]

CELL CYCLE PHASES



2. Human cells divide every day for normal growth, replacement of cells, and tissue repair. The dividing cells proceed through the three phases of the cell cycle as shown in the figure. When a person experiences a cut, damaged skin cells secrete a protein called epidermal growth factor (EGF). Epidermal (skin) cells in the vicinity of the injury possess proteins in their cell membranes called epidermal growth factor receptors (EGFR)
- (a) **Identify** the THREE major stages of interphase, and **describe** the sequence in which they occur.
 - (b) **Describe** the most likely role of EGF and the role of EGFR in a cell signaling pathway. **Describe** how the cellular response to EGF results in tissue repair.
 - (c) Cancer can result from disruptions in cell cycle control. Mutations that increase the production of EGFR have been associated with skin cancer of the head and neck. **Describe** how a mutation outside of the coding region of a gene that encodes EGFR could lead to the overproduction of EGFR protein in a skin cell.
 - (d) In cases of skin cancer that are associated with overproduction of EGFR, treatment with antibodies against the extracellular portion of EGFR has been successful. **Propose** a model to explain how the antibody therapy most likely interferes with the cell signaling pathway, and **give ONE reason** the therapy is effective for treating cancer.

THIS PAGE MAY BE USED FOR TAKING NOTES AND PLANNING YOUR ANSWERS.

NOTES WRITTEN ON THIS PAGE WILL NOT BE SCORED.

WRITE ALL YOUR RESPONSES ON THE LINED PAGES.

GO ON TO THE NEXT PAGE.

[illegible]

[illegible]

[illegible]

[illegible]

DRY MASS OF CORN SEEDLINGS GROWN UNDER DIFFERENT CONDITIONS

Treatment Group	Treatment	Initial Dry Mass of 10 Plants (g)	Dry Mass of 10 Plants After One Week (g)	Change in Dry Mass of 10 Plants Over One Week (g)
I	None	14.8		
II	Light		32.8	+18
III	Dark		11.7	-3.1

3. Thirty corn seedlings of equal size were randomly assigned to one of three treatment groups. At the beginning of the experiment, the plants in group I were dried and the mass was determined. The plants in group II were maintained in light for a week. The plants in group III were maintained in the dark for a week. All conditions, other than light, were the same for groups II and III. At the end of the week, the plants in groups II and III were dried and the mass was determined. The experimental results are provided in the table.
- (a) To explain the increase in mass of the light-grown plants, **identify** ONE inorganic source of new plant mass and **connect** it to the cellular process underlying the increase in mass.
- (b) To explain the decrease in mass of the dark-grown plants, **identify** the overall chemical reaction that is occurring in the plant cells and **connect** it to the cellular process underlying the decrease in mass.

PAGE FOR ANSWERING QUESTION 3

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

[illegible]




Figure 1. Membrane permeability to sodium and potassium ions during an action potential




Figure 2. Membrane potential during an action potential

4. A scientist studying the effect of a stimulus on a motor neuron observed that the permeability of the plasma membrane of the neuron changes during an action potential. The curves in Figure 1 represent the changes in permeability of the membrane to sodium ions (Na^+) and potassium ions (K^+) during the action potential. The graph in Figure 2 represents the changes in membrane potential during an action potential.
- Identify** ONE difference between the Na^+ and K^+ permeability curves shown in Figure 1.
 - Describe** ONE structural feature of the plasma membrane of the motor neuron that regulates the permeability of the plasma membrane to Na^+ and K^+ during an action potential.
 - Connect** the relative membrane permeability data shown in Figure 1 with the changes in membrane potential during stages II and III of the action potential shown in Figure 2.

[illegible]




Figure 1. Blood lactate levels during underwater dives

5. Many aquatic animals can hold their breath and dive underwater for long periods of time. During long dives, lactic acid builds up in tissues of the organism. The graph above shows blood lactate levels following dives of different duration for three species. Lactate is produced when lactic acid is dissolved in the blood and becomes ionized.
- Based on the data, **propose** a hypothesis to explain the change in blood lactate levels in Weddell seals for dives lasting longer than 20 minutes.
 - Describe** the most likely shape of the curve if blood oxygen levels of Weddell seals were plotted rather than blood lactate levels. Include in your description the likely shape of the curve between 0 and 20 minutes and the shape of the curve after 20 minutes.
 - The data suggest that Baikal seals can sustain much longer dives than Emperor penguins. **Propose** a hypothesis that could explain the evolution of different dive responses in Emperor penguins and Baikal seals.

PAGE FOR ANSWERING QUESTION 5

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.




Figure 1. Radioactively labeled polypeptides

6. In an experiment investigating the mechanism of protein secretion, researchers tracked the movement of radioactively labeled polypeptides in pancreatic cells. At various times after the labeling, samples of the cells were observed using an electron microscope to determine the location of the radioactively labeled polypeptides. Figure 1 summarizes the results at 5 minutes, 15 minutes, and 30 minutes. The dark dots in the figure represent the radioactively labeled polypeptides.
- Using the experimental results, **describe** the pathway that secretory proteins take from their synthesis to their release from the cell.
 - Predict** what the results would be if mRNA were radioactively labeled instead of polypeptides. **Give reasoning** to support your prediction.

PAGE FOR ANSWERING QUESTION 6

[illegible]

7. In the early nineteenth century, Alexander von Humboldt observed that the distribution of plants up a mountainside mirrors the distribution of plants at increasing latitude. A number of modern studies have reinforced the idea that as the global climate warms, plant species are becoming redistributed to higher elevations or latitudes.

However, in a separate large-scale study comparing historical (1930–1935) plant distributions with modern (2000–2005) distributions in the western United States, researchers observed that 72% of identified plant species shifted to lower elevations despite an associated increase in average temperature.


Explain the observed shift of the plants to lower elevations by doing each of the following.

- **Propose a hypothesis** about an environmental factor other than temperature that could be driving the shift to lower elevations.
- **Describe** ONE piece of evidence that a researcher could use to support your hypothesis.
- **Provide reasoning** to connect the evidence with the environmental factor you proposed.

PAGE FOR ANSWERING QUESTION 7

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

[illegible]

[illegible]

[illegible]

AP® BIOLOGY EQUATIONS AND FORMULAS

Statistical Analysis and Probability

Mean

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

Standard Deviation

$$s = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n - 1}}$$

Standard Error of the Mean

$$SE_{\bar{x}} = \frac{s}{\sqrt{n}}$$

Chi-Square

$$\chi^2 = \sum \frac{(o - e)^2}{e}$$

Chi-Square Table

p value	Degrees of Freedom							
	1	2	3	4	5	6	7	8
0.05	3.84	5.99	7.82	9.49	11.07	12.59	14.07	15.51
0.01	6.64	9.21	11.34	13.28	15.09	16.81	18.48	20.09

Laws of Probability

If A and B are mutually exclusive, then:

$$P(A \text{ or } B) = P(A) + P(B)$$

If A and B are independent, then:

$$P(A \text{ and } B) = P(A) \times P(B)$$

Hardy-Weinberg Equations

$$p^2 + 2pq + q^2 = 1 \quad p = \text{frequency of the dominant allele in a population}$$

$$p + q = 1 \quad q = \text{frequency of the recessive allele in a population}$$

$\bar{x}$ = sample mean

n = size of the sample

s = sample standard deviation (i.e., the sample-based estimate of the standard deviation of the population)

o = observed results

e = expected results

Degrees of freedom are equal to the number of distinct possible outcomes minus one.

Metric Prefixes

<u>Factor</u>	<u>Prefix</u>	<u>Symbol</u>
10^9	giga	G
10^6	mega	M
10^3	kilo	k
10^{-2}	centi	c
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p

Mode = value that occurs most frequently in a data set

Median = middle value that separates the greater and lesser halves of a data set

Mean = sum of all data points divided by number of data points

Range = value obtained by subtracting the smallest observation (sample minimum) from the greatest (sample maximum)

<p style="text-align: center;">Rate and Growth</p> <p><u>Rate</u> $\frac{dY}{dt}$</p> <p><u>Population Growth</u> $\frac{dN}{dt} = B - D$</p> <p><u>Exponential Growth</u> $\frac{dN}{dt} = r_{\max} N$</p> <p><u>Logistic Growth</u> $\frac{dN}{dt} = r_{\max} N \left(\frac{K - N}{K} \right)$</p> <p><u>Temperature Coefficient Q_{10}</u> $Q_{10} = \left(\frac{k_2}{k_1} \right)^{\frac{10}{T_2 - T_1}}$</p> <p><u>Primary Productivity Calculation</u> $\frac{\text{mg O}_2}{\text{L}} \times \frac{0.698 \text{ mL}}{\text{mg}} = \frac{\text{mL O}_2}{\text{L}}$ $\frac{\text{mL O}_2}{\text{L}} \times \frac{0.536 \text{ mg C fixed}}{\text{mL O}_2} = \frac{\text{mg C fixed}}{\text{L}}$ (at standard temperature and pressure)</p>	<p>dY = amount of change dt = change in time B = birth rate D = death rate N = population size K = carrying capacity $r_{\max}$ = maximum per capita growth rate of population</p> <p>T_2 = higher temperature T_1 = lower temperature k_2 = reaction rate at T_2 k_1 = reaction rate at T_1 Q_{10} = the factor by which the reaction rate increases when the temperature is raised by ten degrees</p>	<p>Water Potential (Ψ) $\Psi = \Psi_P + \Psi_S$ Ψ_P = pressure potential Ψ_S = solute potential</p> <p>The water potential will be equal to the solute potential of a solution in an open container because the pressure potential of the solution in an open container is zero.</p> <p>The Solute Potential of a Solution $\Psi_S = -iCRT$ i = ionization constant (this is 1.0 for sucrose because sucrose does not ionize in water) C = molar concentration R = pressure constant ($R = 0.0831$ liter bars/mole K) T = temperature in Kelvin ($^{\circ}\text{C} + 273$)</p>
<p style="text-align: center;">Surface Area and Volume</p> <p><u>Volume of a Sphere</u> $V = \frac{4}{3} \pi r^3$</p> <p><u>Volume of a Rectangular Solid</u> $V = \ell wh$</p> <p><u>Volume of a Right Cylinder</u> $V = \pi r^2 h$</p> <p><u>Surface Area of a Sphere</u> $A = 4\pi r^2$</p> <p><u>Surface Area of a Cube</u> $A = 6s^2$</p> <p><u>Surface Area of a Rectangular Solid</u> $A = \sum \text{surface area of each side}$</p>	<p>r = radius ℓ = length h = height w = width s = length of one side of a cube A = surface area V = volume Σ = sum of all</p>	<p>Dilution (used to create a dilute solution from a concentrated stock solution) $C_i V_i = C_f V_f$ i = initial (starting) C = concentration of solute f = final (desired) V = volume of solution</p> <p>Gibbs Free Energy $\Delta G = \Delta H - T\Delta S$ ΔG = change in Gibbs free energy ΔS = change in entropy ΔH = change in enthalpy T = absolute temperature (in Kelvin)</p> <p>$\text{pH} = -\log_{10} [\text{H}^+]$</p>

APPENDIX - ANSWERS

AP BIOLOGY 2015

Multiple-choice answer key transcribed from the matching supplied source.

Question	Answer	Question	Answer	Question	Answer
1	B	19	B	37	A
2	D	20	D	38	B
3	C	21	D	39	B
4	C	22	A	40	B
5	A	23	C	41	B
6	C	24	C	42	A
7	D	25	B	43	A
8	A	26	B	44	B
9	B	27	D	45	C
10	A	28	D	46	A
11	B	29	*	47	B
12	C	30	A	48	A
13	B	31	B	49	C
14	B	32	C	50	A
15	C	33	A	51	B
16	C	34	A	52	C
17	B	35	B	53	A
18	C	36	C		

Grid-in answers

121: 40-44

122: 0.12-0.17

123: 13.5

124: 9/16, .5625, .56, .562, .563

125: 4.9-5.1

* Question 29 was not scored in the supplied source; its original status is retained.

AP[®] BIOLOGY
2015 SCORING GUIDELINES

Question 1

To determine the likelihood that a patient with a bacterial infection who is in the intensive care unit (ICU) of a hospital could be successfully treated with antibiotic therapies, researchers investigated the frequency of antibiotic resistance in bacteria isolated from patients in ICUs in the United States from 1994 to 2000. The data from the study are shown in the table, along with the overall change in the frequency of bacterial resistance in patient samples from 1994 to 2000.

INCIDENCE OF ANTIBIOTIC RESISTANCE IN PATIENTS ADMITTED TO ICUs FROM 1994 TO 2000

Antibiotic	Percent of All Samples Demonstrating Antibiotic Resistance in 2000	Overall Change in Percent Resistance to Antibiotics from 1994 to 2000
Ciprofloxacin	19	+10
Gentamicin	22	+6
Ampicillin	65	+3
Imipenem	11	+1
Tazobactam	22	0

- (a) A patient was admitted to the ICU in December 2000 with an uncharacterized bacterial infection. Based on the data from the study **identify** the following.
- The antibiotic that was most likely to be effective for treating the infection.
 - The antibiotic that was least likely to be effective for treating the infection.

Identification (2 points)

- Imipenem is the most effective
- Ampicillin is the least effective

- (b) Using the data, **provide justification** for the claim that antibiotic resistance may arise in bacterial species.

Justification (1 point)

- Resistance to antibiotic(s) in the population is increasing for most of the antibiotics as shown by the overall change in percent resistance to antibiotics from 1994 to 2000.

- (c) **Describe** THREE different processes by which antibiotic resistance develops in individual bacterial cells.

Description (3 points maximum)

- Random mutations lead to changes in DNA sequences
- Transformation is uptake of DNA from the environment to the bacteria
- Conjugation transfers DNA from one bacterium to another during mating
- Transduction introduces DNA to bacteria via viruses
- Horizontal gene transfer introduces new genes to the bacteria (*only if student has not earned a point for transformation, conjugation, or transduction)

AP[®] BIOLOGY
2015 SCORING GUIDELINES

Question 1 (continued)

- (d) **Describe** TWO features of natural selection that lead to the spread of antibiotic resistance in a population over many generations. **Give ONE reason** that the development of antibiotic resistance is an example of evolution.

	Description (1 point per row, 2 points max)
Selective Advantage	<ul style="list-style-type: none">• Variants with antibiotic resistance have higher survival rates• Variants without antibiotic resistance have lower survival rates
Reproductive Fitness	<ul style="list-style-type: none">• Resistant individuals produce more offspring

Reasoning (1 point)

- Phenotypes/characteristics/traits in the population are changing over multiple generations
 - Change in allele frequency in the population
- (e) Giving low doses of antibiotics to beef cattle to control the population sizes of intestinal bacteria can result in more efficient uptake of food materials by the cattle. **Pose** an ethical question that the Food and Drug Administration (FDA) should consider when deciding whether to approve this type of antibiotic use in beef cattle.

Ethical question (1 point)

Question focus may include:


- Harm to humans; e.g. is it appropriate to give low doses of antibiotics to beef cattle if it causes harm to humans?
- Harm to cows; e.g. should we give low doses of antibiotics to beef cattle if it causes harm to the cows?
- Harm to society; e.g. is it okay to give low doses of antibiotics to beef cattle at the risk of spreading antibiotic resistance, and thereby making it more difficult to treat disease?
- Harm to environment; e.g. is it ethical to give low doses of antibiotic to beef cattle if the antibiotics kill environmental microorganisms, and thus disrupt ecosystem processes?

AP[®] BIOLOGY

2015 SCORING GUIDELINES

Question 2

CELL CYCLE PHASES



Human cells divide every day for normal growth, replacement of cells, and tissue repair. The dividing cells proceed through the three phases of the cell cycle as shown in the figure. When a person experiences a cut, damaged skin cells secrete a protein called epidermal growth factor (EGF). Epidermal (skin) cells in the vicinity of the injury possess proteins in their cell membranes called epidermal growth factor receptors (EGFR).

- (a) **Identify** the THREE major stages of interphase, and **describe** the sequence in which they occur.

Identification (3 points maximum)

- Growth/G1
- DNA Synthesis/S
- Preparation for mitosis/G2

Description (1 point)

- Growth, DNA synthesis, preparation for mitosis/G1, S, G2

- (b) **Describe** the most likely role of EGF and the role of EGFR in a cell signaling pathway. **Describe** how the cellular response to EGF results in tissue repair.

Description (3 points maximum)

- EGF is a ligand/binds to EGFR/activates the receptor
- Activated receptor (EGFR) initiates cell signaling pathway/internal signaling cascade
- Cellular response initiates cell cycle progression/cell division

AP[®] BIOLOGY
2015 SCORING GUIDELINES

Question 2 (continued)

- (c) Cancer can result from disruptions in cell cycle control. Mutations that increase the production of EGFR have been associated with skin cancer of the head and neck. **Describe** how a mutation outside of the coding region of a gene that encodes EGFR could lead to the overproduction of EGFR protein in a skin cell.

Description (1 point)

- Mutation in regulatory sequence (promoter or enhancer) could increase rate of transcription of EGFR gene
 - Mutation in transcription factor could increase production of mRNA, which would lead to production of more polypeptide
 - Gene duplication may increase production of EGFR
 - Gene translocation may inactivate repressor
 - Mutation in repressor gene may remove controls on cell division
 - Change in mRNA sequence may increase translation efficiency or mRNA stability
- (d) In cases of skin cancer that are associated with overproduction of EGFR, treatment with antibodies against the extracellular portion of EGFR has been successful. **Propose** a model to explain how the antibody therapy most likely interferes with the cell signaling pathway, and **give ONE reason** the therapy is effective for treating cancer.

Proposed model (1 point)

- Antibody competes with/prevents ligand binding to EGFR
- Receptor activation is inhibited.

Reasoning (1 point)

- Inhibits cell cycle progression
- Blocks cell division

AP[®] BIOLOGY

2015 SCORING GUIDELINES

Question 3

DRY MASS OF CORN SEEDLINGS GROWN UNDER DIFFERENT CONDITIONS

Treatment Group	Treatment	Initial Dry Mass of 10 Plants (g)	Dry Mass of 10 Plants after One Week (g)	Change in Dry Mass of 10 Plants over One Week (g)
I	None	14.8		
II	Light		32.8	+18
III	Dark		11.7	−3.1

Thirty corn seedlings of equal size were randomly assigned to one of three treatment groups. At the beginning of the experiment, the plants in group I were dried and the mass was determined. The plants in group II were maintained in light for a week. The plants in group III were maintained in the dark for a week. All conditions, other than light, were the same for groups II and III. At the end of the week, the plants in groups II and III were dried and the mass was determined. The experimental results are provided in the table.

- (a) To explain the increase in mass of the light-grown plants, **identify** ONE inorganic source of new plant mass and **connect** it to the cellular process underlying the increase in mass.

Identification (1 point)	Connection (1 point)
CO ₂	CO ₂ incorporated into carbohydrates by photosynthesis
H ₂ O	H ₂ O incorporated into carbohydrates by photosynthesis

- (b) To explain the decrease in mass of the dark-grown plants, **identify** the overall chemical reaction that is occurring in the plant cells and **connect** it to the cellular process underlying the decrease in mass.

Identification (1 point)	Connection (1 point)
$\text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 \rightarrow 6\text{CO}_2 + 6\text{H}_2\text{O}$ sugar + oxygen → carbon dioxide + water	CO ₂ released by cellular respiration

AP[®] BIOLOGY
2015 SCORING GUIDELINES

Question 4




Figure 1. Membrane permeability to sodium and potassium ions during an action potential




Figure 2. Membrane potential during an action potential

A scientist studying the effect of a stimulus on a motor neuron observed that the permeability of the plasma membrane of the neuron changes during an action potential. The curves in Figure 1 represent the changes in permeability of the membrane to sodium ions (Na^+) and potassium ions (K^+) during the action potential. The graph in Figure 2 represents the changes in membrane potential during an action potential.

AP[®] BIOLOGY
2015 SCORING GUIDELINES

Question 4 (continued)

- (a) **Identify** ONE difference between the Na^+ and K^+ permeability curves shown in figure 1.

Identification (1 point)

- Peak membrane permeability to Na^+ precedes that of K^+
- Changes in membrane permeability to Na^+ are more rapid than those of K^+
- Changes in membrane permeability to Na^+ are of greater magnitude than those of K^+

- (b) **Describe** ONE structural feature of the plasma membrane of the motor neuron that regulates the permeability of the plasma membrane to Na^+ and K^+ during an action potential.

Description (1 point)

- (Voltage-) Gated channels regulate Na^+ and K^+ movements across membranes
- Phospholipid bilayer prevents unregulated diffusion of Na^+/K^+ across membranes
- Na^+/K^+ pumps maintain membrane potential

- (c) **Connect** the relative membrane permeability data shown in Figure 1 with the changes in membrane potential during stages II and III of the action potential shown in Figure 2.

Stage	Connection between permeability change and membrane potential (1 point per box; 2 points maximum)
II	<ul style="list-style-type: none">• Increased permeability to Na^+ causes depolarization (cell becomes more positive)
III	<ul style="list-style-type: none">• Increased permeability to K^+ causes repolarization (cell becomes more negative)• Decreased permeability to Na^+ causes repolarization (cell becomes more negative)

AP[®] BIOLOGY 2015 SCORING GUIDELINES

Question 5




Figure 1. Blood lactate levels during underwater dives

Many aquatic animals can hold their breath and dive underwater for long periods of time. During long dives, lactic acid builds up in tissues of the organism. The graph above shows blood lactate levels following dives of different duration for three species. Lactate is produced when lactic acid is dissolved in the blood and becomes ionized.

- (a) Based on the data, **propose** a hypothesis to explain the change in blood lactate levels in Weddell seals for dives lasting longer than 20 minutes.

Proposal (1 point)

- Increase in lactate levels are due to fermentation/anaerobic metabolism

- (b) **Describe** the most likely shape of the curve if blood oxygen levels of Weddell seals were plotted rather than blood lactate levels. Include in your description the likely shape of the curve between 0 and 20 minutes and the shape of the curve after 20 minutes.

Description (2 points maximum; points may be earned from only one row)

0-20 min (1 point)	After 20 min (1 point)
Oxygen levels start high and decline steadily	Oxygen levels decline more slowly or remain flat
Oxygen levels start and remain high	Oxygen levels decline

- (c) The data suggest that Baikal seals can sustain much longer dives than Emperor penguins. **Propose** a hypothesis that could explain the evolution of different dive responses in Emperor penguins and Baikal seals.

Proposal (1 point)

- The (genetic/heritable) capacity to sustain dives for longer periods of time provides selective/reproductive advantages (access to food/avoidance of predators) for seals but not for penguins.

AP[®] BIOLOGY 2015 SCORING GUIDELINES

Question 6




Figure 1. Radioactively labeled polypeptides

In an experiment investigating the mechanism of protein secretion, researchers tracked the movement of radioactively labeled polypeptides in pancreatic cells. At various times after the labeling, samples of the cells were observed using an electron microscope to determine the location of the radioactively labeled polypeptides. Figure 1 summarizes the results at 5 minutes, 15 minutes, and 30 minutes. The dark dots in the figure represent the radioactively labeled polypeptides.

- (a) Using the experimental results, **describe** the pathway that secretory proteins take from their synthesis to their release from the cell.

Description (1 point)

(Rough) ER → (vesicle →) Golgi → Vesicle → Exocytosis/fusion with plasma membrane/outside cell

- (b) **Predict** what the results would be if mRNA were radioactively labeled instead of polypeptides.

Give reasoning to support your prediction.

Prediction (1 point)	Reasoning (1 point)
Labeled mRNA is in nucleus first and then moves to cytoplasm/ribosomes/ER (RER)	mRNA is transcribed in nucleus and then moves to cytoplasm for translation

AP[®] BIOLOGY
2015 SCORING GUIDELINES

Question 7

In the early nineteenth century, Alexander von Humboldt observed that the distribution of plants up a mountainside mirrors the distribution of plants at increasing latitude. A number of modern studies have reinforced the idea that as the global climate warms, plant species are becoming redistributed to higher elevations or latitudes.

However, in a separate large-scale study comparing historical (1930–1935) plant distributions with modern (2000–2005) distributions in the western United States, researchers observed that 72% of identified plant species shifted to lower elevations despite an associated increase in average temperature.

Explain the observed shift of the plants to lower elevations by doing each of the following.


- **Propose a hypothesis** about an environmental factor other than temperature that could be driving the shift to lower elevations.
- **Describe** ONE piece of evidence that a researcher could use to support your hypothesis.
- **Provide reasoning** to connect the evidence with the environmental factor you proposed.

3 points maximum (Points may be earned from only one row)

Hypothesis (1 point)	Evidence (1 point) (Must be measurable and express comparison between lower and higher elevations)	Reasoning (1 point)
Water	Higher soil moisture/rainfall data/meteorological data at lower elevations	Low water availability at high elevation limits plant growth
CO ₂	More CO ₂ at low compared to high elevations	Low CO ₂ levels at high elevation limit plant growth
Light (visible and/or UV)	Higher visible light at lower elevations	More photosynthesis with more visible light
	Lower UV light intensity at lower elevations	Less DNA/tissue damage with less UV light
Nutrients	More accessible nutrients in richer soil at lower elevations	Increased nutrient availability can enhance plant growth
Herbivore	More herbivores/herbivory at higher elevations	High herbivory prevents plants from migrating to higher elevations

AP[®] BIOLOGY
2015 SCORING GUIDELINES

Question 8



The graph above shows the initial rate of an enzyme-catalyzed reaction at different substrate concentrations in the presence of a constant concentration of enzyme.

- (a) **Connect** the primary structure of the enzyme to its overall shape.

Connection (1 point)

- The amino acid sequence determines the overall shape (of the polypeptide/protein/enzyme)
- R-groups interact and stabilize the structure

- (b) **Predict** the effect of adding a noncompetitive inhibitor to the reaction mixture on the rate of reaction at a high substrate concentration. **Support** your prediction by describing how a noncompetitive inhibitor affects the structure and function of an enzyme.

Prediction (1 point)

- Noncompetitive/allosteric inhibitor will decrease the (initial) rate of reaction (at any substrate concentration)

Support (1 point)

- Inhibitor binding to site other than active site (allosteric site) changes shape of enzyme, which alters the interaction of substrate with active site