



Mark Scheme (Results)

Summer 2026

Pearson Edexcel International GCSE in Chemistry

4CH1/1CR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at:

www.pearson.com/uk

Summer 2026

Question Paper Log Number P80137A

Publication Code 4CH1_1CR_2606_MS

All the material in this publication is copyright

© Pearson

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a)	neutron		1
(b)	15	ALLOW fifteen	1
(c)	M1 (group) 5 M2 (period) 2		2
(d)	same number of protons and electrons OR each atom has 7 protons and 7 electrons		1
(e) i	nitrogen	ALLOW N	1
ii	3-	ALLOW -3	1
Total for question 1 = 7			

Question number	Answer	Notes	Marks
2 (a)	Oxygen Water	ALLOW air ALLOW moisture/water vapour/steam answer in any order/correct formulae	2
(b) (i)	any one from magnesium/Mg aluminium/Al zinc/Zn		1
(ii)	A lighted/burning/flame splint pops	REJECT a glowing splint IGNORE squeaky pop test alone	1
(c)	D (basic and giant) A is not correct as metals do not form acidic oxides B is not correct as iron(III) oxide does not have a simple structure C is not correct as metals do not form acidic oxides		1
Total for question 2 = 5			


Question number	Answer	Notes	Marks
3	(a) (i) a description including any three of the following points M1 pour (some) solvent into a beaker / chromatography tank M2 (place paper in the solvent/beaker) with the food colourings/spots/(samples of) A, B, C, D are above the level of the solvent OWTTE M3 leave (the paper) until solvent reaches the level shown on the diagram/almost reaches top of paper/when spots reach the level shown on the diagram OWTTE M4 (take the paper out and) leave it to dry	ALLOW any named solvent ALLOW with start/pencil/base line above solvent ALLOW draw line at the level reached/travelled by solvent IGNORE any reference to spot travelling/ any description of the R_f value calculation All marks can be scored /supported from a labelled diagram	3
	(ii) a description containing any two of the following points M1 (food colouring) A contains two dyes M2 (food colouring) A contains (the dye in food colouring) B M3 (food colouring) A contains another dye that is not in any other food colouring (C/D)	ACCEPT compounds/solutes for dyes ALLOW components/ substances/chemicals for dyes IGNORE (food colouring) A is a mixture/ reference to purity/solubility REJECT inks/elements for M1 only ECF for M2 and M3	2
	(iii) use a different solvent OWTTE	ALLOW any named solvent ALLOW thin-layer chromatography IGNORE temperature	1


(b)	<p>M1 distance moved by the dye AND distance moved by the solvent</p> <p>M2 distance moved by the dye $\div$ distance moved by the solvent</p> <p>M3 0.75 – 0.80</p>	<p>answer in the range of 0.75 – 0.80 scores 3</p> <p>ALLOW 4.7 – 4.9 (cm) AND 6.1– 6.3 (cm)</p> <p>ECF from M1</p> <p>ECF from M2 for a correct calculation method only</p> <p>answer to 2 SF or more</p>	3
Total for question 3 = 9			

Question number	Answer		Notes	Marks
4 (a)	Process	Name of change		3
	water changes into ice	freezing		
	solid iodine changes directly into a vapour	sublimation		
	sodium chloride crystals stirred in hot water	dissolving		
(b)	<p>M1 the particles/molecules have more (kinetic) energy</p> <p>M2 so can overcome / break the (intermolecular) forces/forces (between molecules)</p>		<p>IGNORE hot water has more (kinetic) energy</p> <p>ALLOW particles/molecules move faster</p> <p>IGNORE vibrate more</p> <p>ALLOW so can break the bonds (between molecules) OR to break away from one another OR to escape more easily</p> <p>IGNORE references to collisions, activation energy or rate of reaction</p>	2
(c)	<p>M1 measure the melting point</p> <p>M2 the pure solid has a sharp/definite/fixed melting point</p> <p>M3 the mixture/impure solid melts over a range of temperatures</p>		<p>ALLOW measuring freezing point</p> <p>IGNORE reference to boiling point</p> <p>ALLOW the mixture has a lower melting point/a pure substance has a higher melting point</p>	3
Total for question 4 = 8				

Question number	Answer	Notes	Marks
5 (a)	argon carbon dioxide ethene nitrogen		4
(b) (i)	they both have full outer shells (of electrons)	ALLOW they do not need to lose or gain electrons IGNORE 8 electrons alone	1
(ii)	the volume of a gas changes with temperature/ gas expands when hot/heated/gas shrinks when cooled OWTTE	ALLOW to allow the steam to condense	1
(iii)	hydrogen is flammable/explosive	IGNORE temperature control	1
(iv)	 M1 (volume of hydrogen) 157.0-106.7 OR 50.3 cm ³ M2 $(50.3 \div 157.0) \times 100$ OR 32.038% M3 32.0(%) OR M1 $(1 - (106.7 \div 157.0)) = 0.32038$ M2 0.32038×100 M3 32.0(%)	correct answer to 1dp scores 3 ECF for M2 and M3 47.1(%) scores 2 if M1 scored	3
(v)	M1 copper(II) oxide is reduced OR copper(II) oxide loses oxygen OR Cu ²⁺ /copper ion gains electrons M2 hydrogen is oxidised OR hydrogen gains oxygen OR H ₂ /hydrogen loses electrons	REJECT copper alone	2
Total for question 5 = 12			

Question number	Answer	Notes	Marks																
6 (a) (i)	<table border="1"> <tr> <td></td><td>NH_4^+</td><td>Mg^{2+}</td><td>Al^{3+}</td></tr> <tr> <td>Cl^-</td><td>NH_4Cl</td><td>MgCl_2</td><td>AlCl_3</td></tr> <tr> <td>O^{2-}</td><td>no compound formed</td><td>MgO</td><td>Al_2O_3</td></tr> <tr> <td>NO_3^-</td><td>NH_4NO_3</td><td>$\text{Mg}(\text{NO}_3)_2$</td><td>$\text{Al}(\text{NO}_3)_3$</td></tr> </table>		NH_4^+	Mg^{2+}	Al^{3+}	Cl^-	NH_4Cl	MgCl_2	AlCl_3	O^{2-}	no compound formed	MgO	Al_2O_3	NO_3^-	NH_4NO_3	$\text{Mg}(\text{NO}_3)_2$	$\text{Al}(\text{NO}_3)_3$		3
	NH_4^+	Mg^{2+}	Al^{3+}																
Cl^-	NH_4Cl	MgCl_2	AlCl_3																
O^{2-}	no compound formed	MgO	Al_2O_3																
NO_3^-	NH_4NO_3	$\text{Mg}(\text{NO}_3)_2$	$\text{Al}(\text{NO}_3)_3$																
(ii)	ammonium nitrate		1																
(iii)	<p>M1 $(28 \div 80) \times 100$</p> <p>M2 35(g)</p> <p>OR</p> <p>M1 $(100 \div 80 =) 1.25$</p> <p>M2 $1.25 \times 28 = 35 \text{ (g)}$</p>	<p>correct answer scores 2 marks</p> <p>ECF from M1</p> <p>17.5 with/without scores 1</p>	2																
(b) (i)	<p>M1 electrostatic attraction/forces</p> <p>M2 between oppositely charged ions</p>	<p>ACCEPT between cations and anions</p>	2																
(ii)	<p>M1 magnesium loses electrons and chlorine gains electrons</p> <p>M2 one atom of magnesium loses two electrons and two atoms of chlorine each gain one electron</p>	<p>M2 subsumes M1</p> <p>Any mention of sharing of electrons/covalent bonding scores 0</p>	2																
(iii)	<p>M1 add (nitric) acid and silver nitrate</p> <p>M2 white precipitate</p>	<p>REJECT other named acids</p> <p>ALLOW correct formulae</p> <p>M2 dep on silver nitrate in M1</p>	2																
Total for question 6 = 12																			

Question number	Answer	Notes	Marks
7 (a) (i)	all 9 points plotted $\pm$ half a square		1
(ii)	points joined with a curved line of best fit, ignoring the anomalous point at 2 minutes		1
(iii)	the student recorded the volume too early		1
(iv)	tangent drawn to the curve at 4 minutes		1
(v)	M1 suitable values taken from the graph for two coordinates using their tangent	If no tangent drawn then only M3 can be awarded.	3
	 <p>M2 gradient of tangent calculated (expected value = 4.58)</p> <p>M3 cm³/minute</p>	NO ECF from M1	
(b)	<p>M1 the reaction has stopped</p> <p>M2 because all the acid has reacted</p>	<p>ALLOW cm³/min OR cm³min⁻¹</p> <p>IGNORE no more hydrogen/gas produced/reference to the curve plateauing</p>	2
Total for question 7 = 9			

Question number	Answer	Notes	Marks
8 (a) (i)	F		1
(ii)	B		1
(iii)	E		1
(iv)	A		1
(b) (i)	temperature 600-700 °C	ALLOW any temperature in the range ALLOW missing units REJECT any temperature outside the range	2
(ii)	catalyst alumina/silica/aluminosilicates/zeolites M1 shorter alkanes/hydrocarbons (such as C ₈ H ₁₈) are more flammable (than longer alkanes) M2 therefore are more useful/used as fuels M3 (cracking) produces alkenes (such as C ₂ H ₄) M4 which is used to make polymers/plastics	ALLOW correct formulae ALLOW greater demand for short chain alkanes/hydrocarbons (than long chain alkanes) ORA ALLOW short chain alkanes needed for specific uses e.g. petrol/gasoline ALLOW ethanol/alcohol	4
(c)	M1 single bond between carbon 2 and carbon 3 M2 rest of the repeat unit correct  M3 repeat unit in brackets, trailing bonds present and n to the bottom right	REJECT if butane is drawn M3 dep on M2	3
Total for question 8 = 13			

Question number	Answer	Notes	Marks
9 (a)	<p>M1 $((63 \times 62.5) + (65 \times 37.5)) \div 100$ OR $(63 \times 62.5\%) + (65 \times 37.5\%)$</p> <p>M2 63.75 OR 63.8</p>	<p>correct answer scores 2</p> <p>NO ECF from M1</p> <p>Answer of 64 with correct working scores M1</p> <p>Answer of 64 without working scores 0</p>	2
(b) (i)	<p>M1 amount of copper = $158.75 \div 63.5$ OR 2.5 mol</p> <p>M2 A_r of M ($1000 \div 25 =$) 40</p>	M2 subsumes M1	2
(ii)	Ca OR calcium		1
(iii)	<p>M1 M/Ca is the reducing agent</p> <p>M2 M/Ca donates/loses electron(s) (to Cu^{2+})</p>	<p>M2 dep on M1</p> <p>IGNORE M/Ca is oxidised</p>	2
(c)	<p>M1 10 000 mol of CuO needs 5000 mol of C</p> <p>M2 5000 mol of carbon has a mass of 60 kg</p> <p>OR</p> <p>M1 70 kg of carbon = 5833 mol</p> <p>M2 which is enough to react with 11667 mol of CuO</p>	<p>ACCEPT alternative method e.g</p> <p>ALLOW</p> <p>M1 5000 mol of C needed</p> <p>M2 $70000 \div 12 = 5833$ mol used, so C is in excess</p>	2
Total for question 9 = 9			

Question number	Answer	Notes	Marks
10 (a) (i)	neutralisation / acid-base		1
(ii)	(sulfuric) acid gives proton(s)/H ⁺ (to zinc oxide)	<p>ALLOW (sulfuric) acid is a proton donor</p> <p>ALLOW zinc oxide accepts proton(s)</p>	1
(b)	<p>a description that makes reference to the following three points:</p> <p>M1 add zinc carbonate/ powder to acid</p> <p>M2 until no more zinc carbonate reacts</p> <p>M3 filter (off the excess zinc carbonate/powder)</p> <p>AND three of the following points:</p> <p>M4 heat the solution of zinc sulfate until it is partly evaporated/crystals form/saturated solution</p> <p>M5 leave solution to cool/to crystallise</p> <p>M6 filter off crystals</p> <p>M7 suitable method of drying the crystals</p>	<p>ALLOW until there is no more effervescence</p> <p>M3 subsumes M2 as long as excess zinc carbonate is mentioned</p> <p>If heat to dryness only M1, M2, M3 and M4 can be scored</p> <p>If just left the solution to evaporate only M6 and M7 can be scored</p> <p>ALLOW pour off excess liquid/decant off the solution</p> <p>IGNORE references to washing</p> <p>e.g. using filter paper/using paper towel/in a (warm) oven/leave in a warm place to dry</p> <p>REJECT hot oven or any method of direct heating e.g. Bunsen</p> <p>ALLOW leave crystals to dry but not just dry the crystals</p> <p>No M7 if crystals are washed after drying</p>	6

(c)	<p>M1 mass of zinc sulfate = 8.05 (g) AND mass of water = 6.3 (g)</p> <p>M2 amount of zinc sulfate = $8.05 \div 161$ OR 0.05 (mol) AND $6.3 \div 18$ OR 0.35 (mol)</p> <p>M3 $0.35 \div 0.05$ (=7)</p> <p>OR</p> <p>M1 mass of hydrated zinc sulfate = 14.35 (g) AND mass of anhydrous zinc sulfate = 8.05 (g)</p> <p>M2 amount of zinc sulfate = $8.05 \div 161$ OR 0.05 (mol) AND M_r of hydrated zinc sulfate = $(14.35 \div (0.05)) = 287$</p> <p>M3 $(287 - 161) \div 18$ (= 7)</p>	<p>M2 subsumes M1 ECF for M2 only</p> <p>M3 subsumes M1 and M2</p> <p>ALLOW M_r of hydrated zinc sulfate = $14.35 \div (8.05 \div 161) = 287$ for M2</p>	3
(d) (i)	<p>M1 (amount of zinc hydroxide) = $0.250 \div 2$ OR 0.125 (mol)</p> <p>M2 maximum mass of zinc hydroxide = 0.125×99 OR 12.375 (g)</p> <p>M3 $((11.3 \div 12.375) \times 100 =) 91.3(\%)$</p> <p>OR</p> <p>M1 (amount of zinc hydroxide) = $0.250 \div 2$ OR 0.125 (mol)</p> <p>M2 amount of zinc hydroxide produced (= $11.3 \div 99$) = 0.114 (mol)</p> <p>M3 $((0.114 \div 0.125) \times 100 =) 91.3(\%)$</p>	<p>correct answer scores 3</p> <p>M2 subsumes M1</p> <p>ALLOW ECF from M2 ACCEPT 87.8 OR 88(%) if using 0.13×99 in M2</p> <p>ALLOW 45.7(%) scores 2 (step M1 missed)</p> <p>ALLOW any number of SF except 1</p>	3
(ii)	<p>zinc hydroxide/sample is wet/not dry</p> <p>OR</p> <p>contamination of zinc hydroxide/sample is caused by an identified error that would result in a greater mass</p>	<p>ALLOW more ammonium hydroxide added</p> <p>e.g. some left over zinc hydroxide/solids from previous experiments</p>	1
Total for question 10 = 15			

Question number	Answer	Notes	Marks
11 a	they contain one type of atom		1
11 b	M1 (graphite has) weak forces between its layers M2 so layers can slide/slip (over one another)	REJECT weak intermolecular forces/layers of molecules/ions	2
c	diamond M1 giant (covalent) structure/giant lattice M2 strong covalent bonds C ₆₀ fullerene M3 simple molecular (structure) M4 weak intermolecular forces comparison M5 more (thermal/heat) energy is needed to break the covalent bonds (in diamond) than to break / overcome the intermolecular forces (in C ₆₀ fullerene)/to separate the (C ₆₀ fullerene) molecules	ALLOW macromolecular IGNORE giant molecule REJECT ionic ACCEPT strong (electrostatic) forces between the shared pair of electrons and nuclei no M2 if reference to intermolecular forces/ions in diamond ALLOW C ₆₀ exists as molecules ALLOW the attractions between the molecules are weak ALLOW weak intermolecular bonds NOT just heat	5
d	M1 M_r of C ₆₀ = 720 M2 $720 \div 6.02 \times 10^{23}$ M3 mass of 1 molecule = 1.196×10^{-21} (g)	correct answer scores 3 ECF from M1 ECF from M2 for correct calculation method and answer in standard form ALLOW any number of SF except 1	3
total for question 11 = 11			