

INTERNATIONAL AS MATHEMATICS MA02

(9660/MA02) Unit PSM1 Pure Mathematics, Statistics and Mechanics

Mark scheme

January 2026

Version: 1.0 Final

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from www.oxfordaqa.com

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2026 OxfordAQA International Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

M	Mark is for method
m	Mark is dependent on one or more M marks and is for method
A	Mark is dependent on M or m marks and is for accuracy
B	Mark is independent of M or m marks and is for method and accuracy
E	Mark is for explanation
✓ or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
–x EE	Deduct x marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)
ISW	Ignore subsequent working

Q	Answer	Marks	Comments
1	<p>[Perimeter =] $2r + 4.2r [= 124]$</p> <p>[$6.2r = 124 \Rightarrow$] $[r =] 20 \text{ [cm]}$</p> <p>[Area =] $\frac{1}{2} \times 20^2 \times 4.2$</p> <p>[Area =] $840 \text{ [cm}^2\text{]}$</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>oe Correct expression for the perimeter in terms of r PI by the correct value of r or the correct final answer.</p> <p>Correct value of r PI by the correct final answer.</p> <p>oe Correct substitution into $\frac{1}{2}r^2\theta$ ft their r PI by the correct final answer.</p> <p>CAO</p>
		4	
	Question 1 Total	4	

Q	Answer	Marks	Comments
2(a)	$\left[2\sin x + 3\cos x = 0 \right]$ $\Rightarrow \frac{\sin x}{\cos x} = -\frac{3}{2} \Rightarrow$ $\tan x = -\frac{3}{2}$ $[x =] -56.3^\circ, 123.7^\circ$	<p>M1</p> <p>M1</p> <p>A1</p>	<p>oe Deduces the correct value of $\tan x$ Must be identified as $\tan x$ PI by at least one correct final value rounded to the nearest whole number possibly truncated or better ignoring all other values. M1: At least one correct value correctly rounded or both correct values either rounded to the nearest whole number or to 2 or more decimal places. Ignore all other values. A1: CAO</p>
		3	

Q	Answer	Marks	Comments
2(b)	$[5\sin y \cos y] \left(\frac{\cos y}{\sin y} + \frac{\sin y}{\cos y} \right)$ $5\sin y \cos y \left(\frac{\cos^2 y + \sin^2 y}{\sin y \cos y} \right)$ or $5 \left(\frac{\sin y \cos^2 y}{\sin y} + \frac{\cos y \sin^2 y}{\cos y} \right)$ or $\frac{5\sin y \cos^2 y}{\sin y} + \frac{5\cos y \sin^2 y}{\cos y}$ or $5\cos^2 y + 5\sin^2 y$ $5\sin y \cos y \left(\frac{1}{\sin y \cos y} \right) \text{ and } [a =]5$ or $5(\cos^2 y + \sin^2 y) \text{ and } [a =]5$ or $5 \times 1 \text{ and } [a =]5$	<p>M1</p> <p>m1</p> <p>A1</p>	<p>oe Eliminates $\tan y$ using $\tan y = \frac{\sin y}{\cos y}$</p> <p>oe Writes the bracketed term in a full expression with a common denominator or fully or partially expands the bracketed term.</p> <p>Extra line of working showing clear use of $\cos^2 y + \sin^2 y = 1$ before correct value of a given. Must be convincingly shown.</p>
		3	

	Question 2 Total	6	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
3(c)	$[10x^2 - 57x + 77 = 0 \Rightarrow]$ $(5x - 11)(2x - 7) [= 0]$ $[x =] \frac{7}{2} \quad \text{and} \quad [x =] \frac{11}{5}$ $[\text{Area} =]$ $\frac{1}{2} \times \left(\frac{7}{2} + 9\right) \times \left(5 \times \frac{7}{2}\right) \times \frac{\sqrt{51}}{10}$ or $\frac{1}{2} \times \frac{25}{2} \times \frac{35}{2} \times \frac{\sqrt{51}}{10}$ $[\text{Area} =]$ $\frac{175\sqrt{51}}{16} \quad [\text{cm}^2]$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>oe PI by both correct values of $x, \frac{7}{2}$ used to calculate the area, or correct final answer. Correct attempt to solve the quadratic equation Could see use of the quadratic formula but values must be correctly substituted</p> <p>oe Both correct values. PI by $\frac{7}{2}$ used to calculate the area or by the correct final answer.</p> <p>oe Correct use of $\text{Area} = \frac{1}{2}ab \sin C$ with values substituted ft their largest value of x PI by the correct final answer.</p> <p>ACF ISW</p>
		4	
	Question 3 total	9	

Q	Answer	Marks	Comments
4(a)	$\begin{aligned} & \left[x^2 + y^2 = 18x + 4y - 68 \right] \\ & \left[\Rightarrow x^2 - 18x + y^2 - 4y + 68 = 0 \right] \\ & (x-9)^2 \dots \text{ or } (y-2)^2 \dots \\ \\ & (x-9)^2 - 9^2 + (y-2)^2 - 2^2 + 68 [=0] \\ & \text{or} \\ & (x-9)^2 - 81 + (y-2)^2 - 4 + 68 [=0] \\ & \text{or} \\ & (x-9)^2 + (y-2)^2 = 17 \\ \\ & [\text{Centre}] (9,2) \\ \\ & [\text{Radius}] \sqrt{17} \end{aligned}$	<p>M1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>PI by the correct centre.</p> <p>oe Correctly completes both squares simplified or unsimplified. PI by the correct centre and radius.</p> <p>CAO</p> <p>CAO</p>
		4	

Q	Answer	Marks	Comments
4(b)	<p> $[\text{Gradient} =] \frac{3-2}{13-9} \left[= \frac{1}{4} \right]$ </p> <p> $y-3 = -4(x-13)$ or $y-3 = -4x+52$ or $y = -4x+55$ or $4x+y = 55$ </p> <p> $[R] \ (12, 7)$ </p> <p> $[QR ^2 =] (13-12)^2 + (3-7)^2 \ [= 17]$ or $[QR =] \sqrt{(13-12)^2 + (3-7)^2} \ [= \sqrt{17}]$ or $QR ^2 = 17$ or $QR = \sqrt{17}$ </p> <p> $\sqrt{17 - \left(\frac{1}{2}\sqrt{34}\right)^2}$ </p> <p> $\frac{1}{2}\sqrt{34} \quad \text{or} \quad \frac{\sqrt{34}}{2} \quad \text{or} \quad \sqrt{\frac{17}{2}}$ </p>	<p>M1</p> <p>m1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>oe Correct method for finding gradient of radius of C at Q. PI by correct value for the gradient of the radius or the gradient of l_2 ft their centre.</p> <p>oe Correct equation of l_2 ACF ft their centre from part (a) and their gradient of radius</p> <p>Solves equations of l_1 and l_2 simultaneously to find the correct coordinates R</p> <p>oe Correct method to calculate $QR ^2$ or QR or 17 or $\sqrt{17}$ clearly identified. PI by correct final answer. Must be using the correct coordinates of R</p> <p>oe Correct method to find the correct distance to the chord.</p> <p>CAO Correct shortest distance from R to the chord</p>
		6	

Q	Answer	Marks	Comments
4(b) Alt	$[\text{Gradient} =] \frac{3-2}{13-9} \left[= \frac{1}{4} \right]$ <p> $y-3 = -4(x-13)$ or $y-3 = -4x+52$ or $y = -4x+55$ or $4x+y = 55$ </p> <p>$[R] \quad (12, 7)$</p> <p>$[P] \quad (8, 6)$</p> <p> $[\text{Length of } MR =]$ $\sqrt{\left(12-\frac{21}{2}\right)^2 + \left(7-\frac{9}{2}\right)^2}$ </p> <p> $\frac{1}{2}\sqrt{34} \quad \text{or} \quad \frac{\sqrt{34}}{2} \quad \text{or} \quad \sqrt{\frac{17}{2}}$ </p>	<p>M1</p> <p>m1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p>	<p>oe Correct method for finding gradient of radius of C at Q PI by correct value for the gradient of the radius or the gradient of l_2 ft their centre.</p> <p>oe Correct equation of l_2 ACF ft their centre from part (a) and their gradient of radius</p> <p>Solves equations of l_1 and l_2 simultaneously to find the correct coordinates R</p> <p>PI Solves equations of l_1 and C simultaneously to find the correct coordinates P</p> <p>oe Correct method to find the correct distance from R to the midpoint $M\left(\frac{21}{2}, \frac{9}{2}\right)$ of PQ Must be using the correct coordinates of R and M</p> <p>CAO Correct shortest distance from R to the chord</p>
		6	
	Question 4 Total	10	

Q	Answer	Marks	Comments
5	$\left[\frac{9^{5t}}{7^t} = 81 \Rightarrow \frac{9^{5t}}{81} = 7^t \Rightarrow \right]$ $9^{5t-2} = 7^t \quad \text{or} \quad 3^{10t-4} = 7^t$ <p>or</p> $\log_3 9^{5t} - \log_3 7^t = \log_3 81$ <p>or</p> $\log_9 9^{5t} - \log_9 7^t = \log_9 81$ $5t - 2 = \log_9 7^t \quad \text{or} \quad 10t - 4 = \log_3 7^t$ $5t - 2 = t \log_9 7 \quad \text{or} \quad 10t - 4 = t \log_3 7$ $t = \frac{4}{10 - \log_3 7}$	<p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>oe Uses index rules to form a correct equation with $\frac{9^{5t}}{81}$ written as a single power of 9 or 3</p> <p>or</p> <p>oe Takes logarithms of both sides and uses a logarithm rule.</p> <p>oe Takes $\log_9$ or $\log_3$ of both sides to correctly eliminate the power of 9 or 3</p> <p>oe Uses power rule for logarithms to write as a correct linear equation.</p> <p>CAO Must be in the correct form</p>
		4	

	Question 5 Total	4	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
6(a)	$[d =] \log_{10} \left(\frac{b^5}{c} \right) - \log_{10} \left(\frac{b^5}{\sqrt{c}} \right)$ $[d =] \log_{10} \left(\frac{\frac{b^5}{c}}{\frac{b^5}{\sqrt{c}}} \right)$ or $[d =] \log_{10} (b^5) - \log_{10} c - \log_{10} (b^5) + \log_{10} \sqrt{c}$ $[d =] -\log_{10} c + \frac{1}{2} \log_{10} c$ or $[d =] \log_{10} \left(\frac{1}{\sqrt{c}} \right) \text{ or } \log_{10} c^{-\frac{1}{2}}$ or $\frac{1}{2} \log_{10} \left(\frac{1}{c} \right)$ and $[d =] -\frac{1}{2} \log_{10} c$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>oe Correct expression for the common difference in terms of logarithms.</p> <p>Correct expression as a single logarithm or four separate logarithms, for example</p> $\log_{10} \left(\frac{\frac{1}{c}}{\frac{1}{\sqrt{c}}} \right) \text{ or } \log_{10} \left(\frac{\sqrt{c}}{c} \right)$ <p>Must see this extra line of working before AG Must be convincingly shown.</p>
		3	

Q	Answer	Marks	Comments
6(b)	$\frac{1}{2}(15)\left(2\log_{10}\left(\frac{b^5}{\sqrt{c}}\right)+(15-1)\left(-\frac{1}{2}\log_{10} c\right)\right)$ or $\frac{15}{2}\left(2\log_{10}\left(\frac{b^5}{\sqrt{c}}\right)-7\log_{10} c\right)$	M1	oe Correct expression for the sum of the first 15 terms
	$[S_{15} =] \frac{15}{2}\left(\log_{10}\left(\frac{b^{10}}{c}\right)-\log_{10} c^7\right)$	M1	oe Correct application of the power rule for logarithms to both logarithmic terms in a correct expression.
	$[S_{15} =] \frac{15}{2}\log_{10}\left(\frac{b^{10}}{c^8}\right)$	M1	oe Correct application of subtraction rule for logarithms in a correct expression.
	$[S_{15} =] \log_{10}\left(\frac{b^{75}}{c^{60}}\right)$	A1	CAO Must be in the correct form
		4	

	Question 6 Total	7	
--	------------------	---	--

Q	Answer	Marks	Comments
7(a)	$\left[\binom{14}{5} \times 0.35^5 \times (1-0.35)^{14-5} = \right] 0.2178$	B1	AWRT 0.2178
		1	

Q	Answer	Marks	Comments
7(b)	$[P(X > 6) =] 1 - P(X \leq 6)$ or $[P(X > 6) =] 1 - 0.8164$ 0.184	M1 A1	Identifies the correct calculation PI AWRT 0.184
		2	

Q	Answer	Marks	Comments
7(c)	$[P(2 \leq X < 8) =] P(X \leq 7) - P(X \leq 1)$ or $[P(2 \leq X < 8) =] 0.9247 - 0.0205$ 0.904	M1 A1	Identifies the correct calculation PI AWRT 0.904
		2	

	Question 7 Total	5	
--	------------------	---	--

Q	Answer				Marks	Comments								
8(a)	<table><tr><td>x</td><td>0</td><td>100</td><td>200</td></tr><tr><td>$P(X = x)$</td><td>0.25</td><td>0.4125</td><td>0.3375</td></tr></table>				x	0	100	200	$P(X = x)$	0.25	0.4125	0.3375	M1	Correct calculation of one probability
	x	0	100	200										
$P(X = x)$	0.25	0.4125	0.3375											
					A1	Correct calculation of all probabilities								
					2									

Q	Answer	Marks	Comments
8(b)	Mean = $0 \times 0.25 + 100 \times 0.4125 + 200 \times 0.3375$	M1	oe Correct calculation ft their probability distribution
	Mean = 108.75	A1	CAO
		2	

Q	Answer	Marks	Comments
8(c)	$[E(Y) =] \quad 27$	B1	Correct value of $E(Y)$ PI by a final answer of their $E(X)$ minus 27.
	$[E(X - Y) =] \quad 108.75 - 27$	M1	Applies $E(X - Y) = E(X) - E(Y)$ ft their $E(X)$ and $E(Y)$
	$[E(X - Y) =] \quad 81.75$	A1ft	ft their $E(X)$ and $E(Y)$
		3	

	Question 8 Total	7	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
9(a)	$0.6 = k + 2k$	M1	Uses $P(A \cap B) = 0$ and the addition law to form an equation in terms of k oe PI
	$[k =] \ 0.2$	A1	CAO
		2	

Q	Answer	Marks	Comments
9(b)	$[P(A \cap B) =] \ 2k^2$	B1	Uses independence law to find $P(A \cap B)$ in terms of k PI
	$0.6 = k + 2k - 2k^2$	M1	Uses the addition law to form a correct equation in terms of k oe
	$[2k^2 - 3k + 0.6 = 0]$	A1	PI by both possible roots or the correct final answer.
	$[k =] \ 0.238$		AWRT 0.238 If the other root [1.262] is found, it must be rejected
		3	

Q	Answer	Marks	Comments
9(c)	$[P(A \cap B) =] \ k + 2k - 0.6$	B1	Uses addition law to find $P(A \cap B)$ in terms of k May be found in earlier parts but must be used in part (c) to score oe
	$0.5k = \frac{3k - 0.6}{2k}$	M1	PI
	$[k^2 - 3k + 0.6 = 0]$	A1	oe Forms a correct equation in terms of k PI by both possible roots or the correct final answer.
	$[k =] \ 0.215$		Allow 0.2155 or AWRT 0.215 If the other root [2.785] is found, it must be rejected
		3	

	Question 9 Total	8	
--	------------------	---	--

	Answer	Marks	Comments
10(a)	$250 - mg = 0$ or $250 = mg$	M1	Allow inequality signs. PI by correct final answer.
	$[m_{\min} =] 26$	A1	oe AWRT 26 Allow $\frac{1250}{49}$, 25.510204...
		2	

Q	Answer	Marks	Comments
10(b)(i)	$250 - 25g = 25a$ $a = 0.2$	B1	Allow 9.8 for g AG
		1	

Q	Answer	Marks	Comments
10(b)(ii)	$[v = 0 + 0.2 \times 3 =] 0.6$ $[s =] 0 + \frac{1}{2} \times 0.2 \times 3^2 [= 0.9]$ $(0)^2 = (0.6)^2 + 2 \times (-9.8) \times x$ $x = [0.018367\dots]$ or $(0)^2 = (0.6)^2 + 2 \times (-9.8) \times (d - 0.9)$ $[d = 0.9 + 0.018[3\dots] =] 0.92$	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>Achieves velocity as 0.6 PI by 0.6 embedded in equation for motion under gravity or in later working.</p> <p>oe PI by 0.9 seen.</p> <p>oe Sets up an equation(s) to find additional height. Allow sign errors</p> $\left[x = \frac{9}{490} \right] \left[d = \frac{45}{49} \right]$ <p>ft their 0.9 if total height used PI</p> <p>CAO AWRT 0.92 $\left[d = \frac{45}{49} \right]$</p> <p>Their additional height (0.02 or better) must be correctly obtained</p>
		4	
	Question 10 Total	7	

[illegible]

Q	Answer	Marks	Comments
11(b)	Air resistance [not included].	E1	Do not allow reasons based on any values given in the question. Allow 'other resistive forces [not included]'.
		1	

	Question 11 Total	6	
--	--------------------------	----------	--

Q	Answer	Marks	Comments
12(a)	$[v = t^3 - kt^2]$	M1	Correct derivative.
	$\left[\frac{dv}{dt} = \right] 3t^2 - 2kt$		
	$[0.25 = 3k^2 - 2k \times k]$ $k^2 = 0.25]$ $[k =] 0.5$	A1	CAO Must be positive
		2	

Q	Answer	Marks	Comments
12(b)	$[v = t^3 - kt^2] \left[\int v \, dt = \right] \frac{t^4}{4} - \frac{kt^3}{3}$	M1	Correct expression for displacement – allow +C
	$\left[\begin{array}{l} \frac{t^4}{4} - \frac{kt^3}{3} = 0 \\ 3t^4 - 4(0.5)t^3 = 0 \Rightarrow t = \frac{2}{3} \\ 3t - 2 = 0 \end{array} \right]$	A1	Uses their expression for displacement and value for k to find the value t when the particle returns to O PI by $T = \frac{4}{3}k$
	$[0 = t^2(t - k)] \quad [t =] \quad k \quad \text{or} \quad 0.5$	M1	Uses $v = 0$ to find the time for the furthest distance from A
	$\frac{(0.5)^4}{4} - \frac{k(0.5)^3}{3} \quad \left[= -\frac{1}{192} \right]$	M1	Correct method for finding the displacement at $t = k$, Ignore signs for this mark. PI by $-\frac{1}{192}$ or $\frac{1}{96}$ or $\frac{1}{6}k^4$ or $\frac{k^4}{12}$ ft their k provided that it is seen in working
	$\left[\text{Av.Speed} = 2 \times \frac{\left(\frac{1}{192}\right)}{\left(\frac{2}{3}\right)} = \right] \frac{1}{64}$ $[= 0.015625]$	A1	Final answer must be positive Allow exact values, 2.s.f or better
		5	
	Question 12 Total	7	

