

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname _____

Forename(s) _____

Candidate signature _____

I declare this is my own work.

INTERNATIONAL A-LEVEL MATHEMATICS

(9660/MA05) Unit M2 Mechanics

Thursday 4 June 2026

08:00 UK Time

Time allowed: 1 hour 30 minutes

Materials

- For this paper you must have the OxfordAQA Booklet of Formulae and Statistical Tables (enclosed).
- You may use a graphical calculator.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- The **final** answer to questions requiring the use of calculators should be given to two significant figures, unless stated otherwise.
- Unless stated otherwise, the acceleration due to gravity, g , should be taken as 9.8 m s^{-2}

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- Show all necessary working; otherwise marks may be lost.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
TOTAL	



J U N 2 6 M A 0 5 0 1

IB/M/Jun26/G4006/V4

MA05

Answer **all** questions in the spaces provided.

- 1** Four particles P , Q , R and S are fixed on a uniform rectangular lamina $OABC$ to create a system. The system is shown in **Figure 1**

Figure 1




Figure 2 shows the mass of each particle and the coordinates of the points at which they are fixed to the lamina.

Figure 2

Particle	Mass (kg)	Coordinates (metres)
P	3	(2, 2)
Q	4	(2, 8)
R	7	(10, 8)
S	9	(10, 2)

The mass of the uniform rectangular lamina $OABC$ is 11 kg

The coordinates of the point A are (0, 10)

The coordinates of the point C are (12, 0)

- 1 (a)** State, with a reason, the coordinates of the centre of mass of the **uniform rectangular lamina**.

[2 marks]

Coordinates _____

Reason _____



Give your answer in an exact form.

[illegible]

Give your answer to the nearest degree.

9

2 A particle of mass 4 kg is moving up a smooth inclined plane.

The plane is inclined at an angle θ to the horizontal.

The particle passes through the points X and Y

As the particle passes through X it is moving with speed 3.2 m s^{-1} up the plane.


As the particle passes through Y it is moving with speed 1.2 m s^{-1} up the plane.

The line through X and Y is a line of greatest slope on the inclined plane.

The distance between X and Y is d metres.

The inclined plane is shown in **Figure 3**

Figure 3



Assume that the particle experiences no resistance forces or friction forces.

2 (a) Show that the loss in the particle's kinetic energy between X and Y is 17.6 J

[1 mark]

- 2 (b)** It is given that the particle experiences a normal reaction force of magnitude 25 newtons due to its contact with the plane.

- 2 (b) (i)** Find the value of θ

Give your answer to the nearest degree.

[2 marks]

$\theta =$ _____

- 2 (b) (ii)** Hence use an energy method to find the value of d

[3 marks]

$d =$ _____

- 2 (c)** The particle continues to move up the plane until it momentarily comes to rest.

The particle then begins to move down the plane.

The total time that the particle is between X and Y during its entire motion on the plane is T seconds.

Find the value of T

[2 marks]

$T =$ _____



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



6

- 4** A car of mass 1200 kg moves at constant speed $v \text{ m s}^{-1}$ around a roundabout, which is a circular section of road of radius 50 metres.

The position vector of the car $\mathbf{r}$ metres relative to the centre of the roundabout at time t seconds is given by

$$\mathbf{r} = \begin{bmatrix} 50 \cos(0.24t) \\ 50 \sin(0.24t) \end{bmatrix}$$

The road on the roundabout is horizontal and **dry**.

The coefficient of friction between the car's tyres and the **dry** road is μ

- 4 (a) (i)** Write down the angular speed of the car as it moves around the roundabout.

Include units in your answer.

[2 marks]

Answer _____ units _____

- 4 (a) (ii)** Find the value of v

[2 marks]

$v =$ _____

- 4 (a) (iii)** Find the range of values of μ for which the car can move with constant speed $v \text{ m s}^{-1}$ around the roundabout in a circle of radius 50 metres.

[2 marks]

Answer _____



4 (b) A rain shower makes the road on the roundabout **wet**.

It is given that the coefficient of friction between the car's tyres and the **wet** road is 0.18

State, with a reason, whether the car can still move around the roundabout in a circle of radius 50 metres with the same constant speed $v \text{ m s}^{-1}$

[2 marks]

8

Turn over for the next question

Turn over ►



5

$$\mathbf{F} = \begin{bmatrix} 4e^{2t} \\ 16 \cos 4t \\ -12 \sin 3t \end{bmatrix}$$

Assume that no other forces act on the body.

5 (a)

It is given that the initial velocity of the body is $\begin{bmatrix} 2 \\ -5 \\ 3 \end{bmatrix} \text{ m s}^{-1}$

At time t seconds, the velocity of the body is $\mathbf{v} \text{ m s}^{-1}$

Find $\mathbf{v}$ in terms of t

[6 marks]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Answer



5 (b) Hence find the speed of the body when $t = \frac{\pi}{12}$

[2 marks]

Answer _____

8

Turn over for the next question

Turn over ►



Assume that the only force the ball experiences whilst it is in the air is its weight.

$$\frac{gx^2}{2y^2} \tan^2 \theta - x \tan \theta + y + \frac{gx^2}{2y^2} = 0$$

[6 marks]

[illegible]

*Do not write
outside the
box*


- 6 (b)** A child is trying to kick a small ball from a point O on horizontal ground to make it pass over a vertical wall.

The child kicks the ball so its initial speed as it leaves their foot is 8 m s^{-1}

The horizontal displacement from O to the bottom of the wall is 5 metres.

The top of the wall is 1.5 metres vertically above the ground as shown in **Figure 4**

Figure 4



Use the result in **part (a)** to determine whether the child can kick the ball to make it pass over the wall.

[4 marks]

[illegible]

7

Two rough fixed planes are both inclined at 25° to the horizontal.


A smooth peg is located at P where the planes meet.

Two particles A and B are connected by a light inextensible string which passes over the peg.

Particle A has mass 5 kg and particle B has mass 11 kg

The particles are held at rest with the string taut and parallel to the line of greatest slope of each plane as shown in **Figure 5**

Figure 5



The coefficient of friction between the particles and each plane is μ

Both particles are released at the same time.

Exactly 4 seconds after being released, A has travelled a distance of 3 metres up the plane towards P

7 (a) (i) Find the magnitude of the acceleration of A

[1 mark]

Answer _____



7 (a) (ii) State with a reason the magnitude of the acceleration of B

[2 marks]

Magnitude _____

Reason _____

7 (b) Show that the value of μ is 0.133 correct to three significant figures.

[5 marks]

Question 7 continues on the next page

Turn over ►



Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Turn over ►



The horizontal circle has radius r metres and centre C as shown in **Figure 7**

A diagram showing a circular path represented by a dashed ellipse. A horizontal line segment with arrows at both ends indicates the radius, labeled r metres. The right endpoint of this segment is marked with a dot and labeled C , representing the center of the path. The path itself has arrows indicating a counter-clockwise direction of travel.

$$F = \frac{k}{r^n}$$

[7 marks]


[illegible]

[illegible]

$k =$ _____ $n =$ _____

7

- 9** A uniform rod of mass 20 kg has length 5 metres.
- The rod rests with one end on a smooth inclined plane at the point A
- The plane is inclined at an angle of 15° above the horizontal.
- The other end of the rod rests against a smooth vertical wall at the point B
- The rod is in equilibrium and makes an angle of θ with the smooth vertical wall, as shown in **Figure 8**

Figure 8

- 9 (a)** The magnitude of the reaction force exerted on the rod at A due to its contact with the inclined plane is R newtons.
- By considering the equilibrium of the forces acting on the rod, show that $R = 203$ correct to three significant figures.

[2 marks]



- 9 (b) (i)** Find the magnitude of the reaction force exerted on the rod due to its contact with the wall at B .

Give your value to three significant figures.

[2 marks]

Answer _____

- 9 (b) (ii)** State with a reason the magnitude of the reaction force exerted on the wall at B due to its contact with the rod.

[2 marks]

Magnitude _____

Reason _____

- 9 (c)** Find the angle θ

Give your answer to the nearest degree.

[4 marks]

$\theta =$ _____

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



*Do not write
outside the
box*

[illegible]

*Do not write
outside the
box*

[illegible]