

INTERNATIONAL A-LEVEL FURTHER MATHEMATICS

FM03

(9665/FM03) Unit FP2 Pure Mathematics

Mark scheme

June 2025

Version: 1.0 Final



2 5 6 X F M 0 3 / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from www.oxfordaqa.com

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2025 OxfordAQA International Examinations and its licensors. All rights reserved.

Key to mark scheme abbreviations

M	Mark is for method
m	Mark is dependent on one or more M marks and is for method
A	Mark is dependent on M or m marks and is for accuracy
B	Mark is independent of M or m marks and is for method and accuracy
E	Mark is for explanation
✓ or ft	Follow through from previous incorrect result
CAO	Correct answer only
CSO	Correct solution only
AWFW	Anything which falls within
AWRT	Anything which rounds to
ACF	Any correct form
AG	Answer given
SC	Special case
oe	Or equivalent
A2, 1	2 or 1 (or 0) accuracy marks
–x EE	Deduct x marks for each error
NMS	No method shown
PI	Possibly implied
SCA	Substantially correct approach
sf	Significant figure(s)
dp	Decimal place(s)
ISW	Ignore subsequent working

Q	Answer	Marks	Comments
1	$\overrightarrow{AB} = \begin{bmatrix} 5 \\ 1 \\ 2 \end{bmatrix}, \quad \overrightarrow{AC} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ $\overrightarrow{AB} \times \overrightarrow{AC} = \begin{bmatrix} 5 \\ 1 \\ 2 \end{bmatrix} \times \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -3 \\ 1 \\ 7 \end{bmatrix}$ $\text{Area} = \frac{1}{2} \left\ \begin{bmatrix} -3 \\ 1 \\ 7 \end{bmatrix} \right\ $ $= \frac{\sqrt{59}}{2}$	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>Two correct directional vectors</p> $\overrightarrow{BC} = \begin{bmatrix} -2 \\ 1 \\ -1 \end{bmatrix}$ <p>Cross product of their directional vectors – not position vectors</p> $\overrightarrow{BA} \times \overrightarrow{BC} = \begin{bmatrix} 3 \\ -1 \\ -7 \end{bmatrix}$ $\overrightarrow{CA} \times \overrightarrow{CB} = \begin{bmatrix} -3 \\ 1 \\ 7 \end{bmatrix}$ <p>Uses correct formula for area of a triangle with their cross product</p> <p>Correct exact answer</p>
	Question 1 Total	4	

Q	Answer	Marks	Comments
3(b)	$\frac{-p}{1} = 4 - 3i + 4 + 3i + 5$ $p = -13$ $\frac{-q}{1} = (4 - 3i) \times (4 + 3i) \times 5$ $q = -125$ $p = -13$ $q = -125$	<p>M1</p> <p>Correct use of $\alpha + \beta + \gamma = \frac{-b}{a}$ and $\alpha\beta\gamma = \frac{-d}{a}$ with their roots to obtain values for p and q</p> <p>Condone $\alpha + \beta + \gamma = \pm \frac{b}{a}$ and $\alpha\beta\gamma = \pm \frac{d}{a}$ or</p> <p>Expands $(z^2 - 8z + 25)(z + \gamma)$ and compares coefficients to obtain values for p and q</p> <p>Condone $(z^2 \pm 8z \pm 25)(z + \gamma)$ or</p> <p>Substitutes a complex root into the given equation and expands to obtain two equations (one for the Real part & one for the Imaginary part)</p> <p>A1</p> <p>Obtains the correct values of p and q</p> <p>If 0/2 scored SC1 for writing $p = -(\alpha + \beta + \gamma)$ and $q = -\alpha\beta\gamma$</p>	
		2	
	Question 3 Total	5	

Q	Answer	Marks	Comments
4	$z^4 = 16i = 16e^{\frac{\pi}{2}i}$ $z^4 = 16e^{\left(\frac{\pi}{2} + 2\pi k\right)i}$ $z = 2e^{\frac{\pi(1+4k)}{8}i}$ $r = 2$ $\theta = \frac{\pi}{8}, \frac{5\pi}{8}, \frac{9\pi}{8}, \frac{13\pi}{8}$ $[z =] 2e^{\frac{\pi}{8}i}, 2e^{\frac{5\pi}{8}i}, 2e^{\frac{9\pi}{8}i}, 2e^{\frac{13\pi}{8}i}$	<p>B1</p> <p>M1</p> <p>B1</p> <p>A1</p> <p>A1</p>	<p>Correct Eulerian form of 16i Accept mod arg form</p> <p>Divides their argument by 4</p> <p>Correct value of r</p> <p>Correct angles (mod 2π)</p> <p>Correct with $0 \leq \theta < 2\pi$</p>
	Question 4 Total	5	

Q	Answer	Marks	Comments
5(a)	$\frac{1}{2}(k+1)[4(k+1)^2 + 7(k+1) + 1]$	M1	Expands and collects to get at least one of b , c or d correct
	$= \frac{1}{2}(k+1)(4k^2 + 8k + 4 + 7k + 7 + 1)$		
	$= \frac{1}{2}(k+1)(4k^2 + 15k + 12)$		
	$= \frac{1}{2}(4k^3 + 19k^2 + 27k + 12)$	A1	Correctly obtains the required result
		2	

Q	Answer	Marks	Comments
5(b)	LHS = $2 \times 3 = 6$ RHS = $\frac{1}{2} \times 1 \times (4 \times 1^2 + 7 \times 1 + 1) = 6$ [True for $n = 1$]	B1	Demonstrates result is true for $n = 1$
	$n = k + 1$ Assume true $n = k$ (*) $2 \times 3 + 5 \times 5 + 8 \times 7 + 11 \times 9 + \dots + (3k - 1) \times (2k + 1)$ $+ [3(k + 1) - 1] \times [2(k + 1) + 1]$		
	$= \sum_{r=1}^k (3r - 1) \times (2r + 1) + [3(k + 1) - 1] \times [2(k + 1) + 1]$ $= \frac{1}{2}k(4k^2 + 7k + 1) + [3(k + 1) - 1] \times [2(k + 1) + 1]$	M1	Assumes true for $n = k$ and adds $[3(k + 1) - 1] \times [2(k + 1) + 1]$ to $= \frac{1}{2}k(4k^2 + 7k + 1)$ Condone lack of $2 \times 3 + 5 \times 5 + 8 \times 7 + 11 \times 9 + \dots + (3k - 1) \times (2k + 1)$ $+ [3(k + 1) - 1] \times [2(k + 1) + 1] =$
	$= \frac{1}{2}k(4k^2 + 7k + 1) + [3(k + 1) - 1] \times [2(k + 1) + 1]$ $= \frac{1}{2}(4k^3 + 7k^2 + k) + (6k^2 + 13k + 6)$ $= \frac{1}{2}(4k^3 + 19k^2 + 27k + 12)$	M1	Expands and collects terms

	$= \frac{1}{2}(k+1)\left[4(k+1)^2 + 7(k+1) + 1\right]$ <p>from (a)</p>
--	--

	Question 5 Total	8	
--	------------------	---	--

Q	Answer	Marks	Comments
6	$\frac{dx}{dt} = t^2 - t$ $\frac{dy}{dt} = 2t^{\frac{3}{2}}$ $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = t^4 - 2t^3 + t^2 + 4t^3$ $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = t^4 + 2t^3 + t^2$ $= (t^2 + t)^2$ $PQ = \int_0^{\sqrt{2}} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$ $PQ = \int_0^{\sqrt{2}} \sqrt{(t^2 - t)^2 + (2t^{\frac{3}{2}})^2} dt$ $PQ = \int_0^{\sqrt{2}} (t^2 + t) dt$ $PQ = \left[\frac{t^3}{3} + \frac{t^2}{2} \right]_0^{\sqrt{2}} = \frac{(\sqrt{2})^3}{3} + \frac{(\sqrt{2})^2}{2}$ $PQ = \frac{2\sqrt{2}}{3} + 1$	<p>B1</p> <p>M1</p> <p>A1</p> <p>B1ft</p> <p>M1</p> <p>A1</p>	<p>Correct expression for $\frac{dx}{dt}$ and $\frac{dy}{dt}$</p> <p>Correctly expands their $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2$</p> <p>$\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = (t^2 + t)^2$ oe</p> <p>Their correct integrand (can be unsimplified), with dt and correct limits. All seen here</p> <p>Correct use of limits in an expression of the form $mt^3 + nt^2$</p> <p>CSO – form of answer given B1 M1 A1 B0 M1 A1 is possible</p>
	Question 6 Total	6	

Q	Answer	Marks	Comments
7(a)	$0 = \begin{vmatrix} 2-k & 1 & 2 \\ 1 & 3 & -2 \\ 8 & 4 & -5k-1 \end{vmatrix}$ $= (2-k)\{3(-5k-1)+8\}$ $-(-5k-1+16)+2(4-24)$ $0 = 15k^2 - 35k + 10 + 5k - 55$ $= 15k^2 - 30k - 45$ $= k^2 - 2k - 3$ $k = -1 \text{ or } k = 3$	<p>M1</p> <p>A1</p> <p>A1</p>	<p>Sets determinant = 0 and attempts to expand</p> <p>Obtains a correct quadratic equation</p> <p>Obtains correct values of k</p>
		3	

Q	Answer	Marks	Comments
7(b)(i)	$\begin{aligned} 3x + y + 2z &= 5 & \text{(I)} \\ x + 3y - 2z &= -1 & \text{(II)} \\ 8x + 4y + 4z &= 12 & \text{(III)} \end{aligned}$ $\begin{aligned} \text{(I)} + \text{(II)} &\Rightarrow 4x + 4y = 4 \\ \text{(III)} + 2\text{(II)} &\Rightarrow 10x + 10y = 10 \end{aligned}$ <p>Infinite solutions</p>	<p>B1ft</p> <p>M1</p> <p>A1</p>	<p>Correct system of equations for their smallest k</p> <p>Eliminating one variable in order to compare two simultaneous equations</p> <p>From comparing correct equations</p>
		3	

Q	Answer	Marks	Comments
7(b)(ii)	Consistent	E1ft	Must follow from their answer to part (b)(i)
		1	

Q	Answer	Marks	Comments
7(b)(iii)	Line of intersection	E1	CAO, oe Must come from correct working with correct equations in part (b)(i)
		1	

	Question 7 Total	8	
--	------------------	---	--

Q	Answer	Marks	Comments
8(a)	$\frac{A}{r} + \frac{B}{r+1} + \frac{C}{r+2}$ $4r^2 + 8r + 2 = A(r+1)(r+2) + Br(r+2) + Cr(r+1)$ $r=0: 2=2A; r=-1: -2=-B; r=-2: 2=2C$ $\frac{1}{r} + \frac{2}{r+1} + \frac{1}{r+2}$	<p>M1</p> <p>A1</p>	<p> $4r^2 + 8r + 2 = A(r+1)(r+2)$ $+ Br(r+2) + Cr(r+1)$ used to form either at least two correct equations in A, B or C or two correct simultaneous equations in A, B and C </p> <p>Correct partial fractions</p>
		2	

Q	Answer	Marks	Comments
8(b)	$\sum_{r=1}^{2n} \frac{(-1)^r (4r^2 + 8r + 2)}{r(r+1)(r+2)}$ $= -\frac{1}{1} - \frac{2}{2} - \frac{1}{3}$ $+ \frac{1}{2} + \frac{2}{3} + \frac{1}{4}$ $- \frac{1}{3} - \frac{2}{4} - \frac{1}{5}$ $\vdots$ $+ \frac{1}{2n-2} + \frac{2}{2n-1} + \frac{1}{2n}$ $- \frac{1}{2n-1} - \frac{2}{2n} - \frac{1}{2n+1}$ $+ \frac{1}{2n} + \frac{2}{2n+1} + \frac{1}{2n+2}$ $= -\frac{1}{1} - \frac{1}{2} + \frac{1}{2n+1} + \frac{1}{2n+2}$ $= -\frac{3}{2} + \frac{4n+3}{(2n+1)(2n+2)}$	<p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>Writes at least three lines of subtracting terms, using their part (a)</p> <p>Writes at least three consecutive lines, showing cancellation</p> <p>Correctly reduces the expression to four terms</p> <p>Obtains the required result This mark is only available if at least the first three lines and the last three lines are present. AG</p>
		4	

Q	Answer	Marks	Comments
8(c)	$-\frac{3}{2}$	B1	CAO
		1	

	Question 8 Total	7	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
9(a)(i)	$\left[\frac{d^6 y}{dx^6} = 64 \cosh 2x \right]$	B1	ACF
		1	

Q	Answer	Marks	Comments
9(a)(ii)	<p>When $x = 0$:</p> $y = 1, \quad \frac{dy}{dx} = 0, \quad \frac{d^2 y}{dx^2} = 4, \quad \frac{d^3 y}{dx^3} = 0,$ $\frac{d^4 y}{dx^4} = 16, \quad \frac{d^5 y}{dx^5} = 0, \quad \frac{d^6 y}{dx^6} = 64$ <p>Maclaurin series:</p> $= 1 + x \times 0 + \frac{x^2}{2!} \times 4 + \frac{x^3}{3!} \times 0$ $+ \frac{x^4}{4!} \times 16 + \frac{x^5}{5!} \times 0 + \frac{x^6}{6!} \times 64$ $[\cosh 2x =] 1 + 2x^2 + \frac{2}{3}x^4 + \frac{4}{45}x^6$	<p>B1</p> <p>M1</p> <p>A1</p>	<p>ft Their derivatives</p> <p>AG Must be convincingly shown</p>
		3	

Q	Answer	Marks	Comments
9(b)	$\cos 2x = 1 - 2x^2 + \frac{2}{3}x^4 - \frac{4}{45}x^6 + \dots$ $[\ln(1+x)]^2 = \left[x - \frac{x^2}{2}\right]^2 = x^2 - x^3 + \dots$ $\frac{\cosh 2x - \cos 2x}{[\ln(1+x)]^2} = \frac{4x^2 + \frac{8}{45}x^6 + \dots}{x^2 - x^3 + \dots}$ $\lim_{x \rightarrow 0} \left[\frac{\cosh 2x - \cos 2x}{[\ln(1+x)]^2} \right]$ $= \lim_{x \rightarrow 0} \left[\frac{4x^2 + \frac{8}{45}x^6 + \dots}{x^2 - x^3 + \dots} \right]$ $= \lim_{x \rightarrow 0} \left[\frac{4 + \frac{8}{45}x^4 + \dots}{1 - x + \dots} \right]$ <p>So limit exists</p> <p>= 4</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>E1</p> <p>A1</p>	<p>Condone</p> $\cos 2x = 1 - \frac{4}{2!}x^2 + \frac{16}{4!}x^4 - \frac{64}{6!}x^6$ <p>May be on last line May see $O(x^4)$ and/or $O(x)$</p> <p>With comment</p> <p>B1 B1 B1 M1 E0 A1 is possible</p>
		6	
	Question 9 Total	10	

Q	Answer	Marks	Comments
10(a)	<p>I.F. is $e^{-\int \frac{1}{x} dx} = e^{-\ln(x)}$</p> <p>$= \frac{1}{x}$</p> <p>$\frac{y}{x} = \int \left(\frac{1}{\sqrt{1-x^2}} \sin^{-1}x \right) dx$</p> <p>$u = \sin^{-1}x \quad \frac{dv}{dx} = \frac{1}{\sqrt{1-x^2}}$</p> <p>$\frac{du}{dx} = \frac{1}{\sqrt{1-x^2}} \quad v = \sin^{-1}x$</p> <p>$\int \left(\frac{1}{\sqrt{1-x^2}} \sin^{-1}x \right) dx$</p> <p>$= \left[(\sin^{-1}x)^2 \right] - \int \left(\frac{1}{\sqrt{1-x^2}} \sin^{-1}x \right) dx$</p> <p>$\int \left(\frac{1}{\sqrt{1-x^2}} \sin^{-1}x \right) dx =$</p> <p>$\frac{1}{2} \left[(\sin^{-1}x)^2 \right]$</p> <p>$y = x \left[\frac{1}{2} (\sin^{-1}x)^2 + c \right]$</p>	<p>M1</p> <p>A1</p> <p>m1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>I.F. identified and integration attempted</p> <p>Condone I.F. $= +\frac{1}{x}$</p> <p>Correct integrating factor</p> <p>Multiplying both sides of the Standard Form by their I.F. and integrating LHS to get $y \times \text{I.F.}$</p> <p>Uses Integration by Parts or Uses Integration by Substitution</p> <p>Correct use of Integration by Parts or Correct use of Integration by Substitution</p> <p>Obtains $\frac{1}{2} \left[(\sin^{-1}x)^2 \right]$</p> <p>Correct result ACF, must be $y = \dots$ ISW</p>
		7	

Q	Answer	Marks	Comments
10(b)	$\frac{\pi^2}{4} = \frac{1}{2} \left(\frac{\pi}{2} \right)^2 + c$ $c = \frac{\pi^2}{8}$ $y = x \left[\frac{1}{2} (\sin^{-1} x)^2 + \frac{\pi^2}{8} \right]$	<p>M1</p> <p>A1</p>	<p>Substituting $y = \frac{\pi^2}{4}$ and $x = 1$ into their GS (with a constant of integration) to find a value of c</p> <p>Correct PS ACF, must be $y = \dots$ ISW</p>
		2	
	Question 10 Total	9	

Q	Answer	Marks	Comments
11(a)(i)		<p>M1</p> <p>Correct shape with x intercept clearly > 0 (does not need to be labelled). Graph in first quadrant only.</p> <p>A1</p> <p>Fully correct with correct x intercept labelled (0.5)</p>	
		2	

Q	Answer	Marks	Comments
11(a)(ii)	$\cosh y = 2x$ $\frac{dy}{dx} \sinh y = 2$ $\frac{dy}{dx} = \frac{2}{\sinh y} = \frac{\pm 2}{\sqrt{\cosh^2 y - 1}}$ $\frac{dy}{dx} = \frac{\pm 2}{\sqrt{4x^2 - 1}}$ [From graph] $\frac{dy}{dx} > 0$ $\therefore \frac{dy}{dx} = \frac{2}{\sqrt{4x^2 - 1}}$	<p>B1</p> <p>Correctly differentiates equation</p> <p>M1</p> <p>Use of $\sinh^2 y = \cosh^2 y - 1$ in their differential equation</p> <p>Condone lack of $\pm$</p> <p>A1</p> <p>Correct differential equation including $\pm$</p> <p>E1</p> <p>Correct explanation leading to required result. AG</p>	
		4	

Q	Answer	Marks	Comments
11(b)	$[f'(x) =] -2x + \frac{2}{\sqrt{4x^2 - 1}}$ $-2x + \frac{2}{\sqrt{4x^2 - 1}} = 0$ $x = \frac{1}{\sqrt{4x^2 - 1}}$ $x\sqrt{4x^2 - 1} = 1$ $x^2(4x^2 - 1) = 1$ $4(x^2)^2 - x^2 - 1 = 0$ $x^2 = \frac{1 - \sqrt{17}}{8} \text{ or } x^2 = \frac{1 + \sqrt{17}}{8}$ $x^2 = \frac{1 - \sqrt{17}}{8} < 0 \text{ reject as } x \in \mathbb{R}$ $x^2 = \frac{1 + \sqrt{17}}{8}$ $x = -\sqrt{\frac{1 + \sqrt{17}}{8}} \text{ reject as } x \geq \frac{1}{2}$ $\therefore x = \sqrt{\frac{1 + \sqrt{17}}{8}}$ <p>Hence $f(x)$ has a single stationary point</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>E1</p> <p>E1</p> <p>A1</p>	<p>Correctly differentiates $f(x)$</p> <p>Obtains a quadratic equation in x^2</p> <p>Correct exact roots</p> <p>Correct rejection of $x^2 < 0$ roots Must have correct exact roots</p> <p>Correct rejection of $x = -\sqrt{\frac{1 + \sqrt{17}}{8}}$ root. Must reference $x \geq \frac{1}{2}$ oe Condone $x > \frac{1}{2}$ Must have correct roots</p> <p>ACF A1 with correct conclusion CSO Must have been awarded B1 M1 A1 E1 E1 to obtain this mark</p>
		6	
	Question 11 Total	12	

Q	Answer	Marks	Comments
12(a)	$\mathbf{p} = \overrightarrow{AB} = \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}, \quad \mathbf{q} = \overrightarrow{AC} = \begin{bmatrix} -2 \\ -3 \\ -3 \end{bmatrix}$	M1 A1	M1 : clear attempt to calculate two directional vectors A1 : two correct directional vectors
	$\mathbf{r} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix} + \mu \begin{bmatrix} -2 \\ -3 \\ -3 \end{bmatrix}$	A1ft	oe In correct form
		3	

Q	Answer	Marks	Comments
12(b)	$\mathbf{n}_1 = \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix} \times \begin{bmatrix} -2 \\ -3 \\ -3 \end{bmatrix} = \begin{bmatrix} 6 \\ -5 \\ 1 \end{bmatrix}$	B1ft	Calculates the cross product of their directional vectors from part (a)
	$\mathbf{n}_1 \times \mathbf{n}_2 = \begin{bmatrix} 6 \\ -5 \\ 1 \end{bmatrix} \times \begin{bmatrix} 2 \\ -1 \\ -3 \end{bmatrix}$	M1	Calculates cross product of their normal vectors
	$\mathbf{v} = \alpha \begin{bmatrix} 4 \\ 5 \\ 1 \end{bmatrix}$	A1	Correct v (accept any scalar multiple)
	$d_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \bullet \begin{bmatrix} 6 \\ -5 \\ 1 \end{bmatrix} = 3$	M1	Calculates the scalar product of their $\mathbf{n}_1$ and the position vector of A, B or C or Substitutes in parametric form of Π_1 into equation for Π_2
	$6x - 5y + z = 3$	A1	Correct Cartesian form of Π_1 PI Later working or Correct parametric equation

	$z = 0$ $6x - 5y = 3$ $2x - y = 7$ $x = 8, y = 9$ $\mathbf{u} = \begin{bmatrix} 8 \\ 9 \\ 0 \end{bmatrix}$ $\left(\mathbf{r} - \begin{bmatrix} 8 \\ 9 \\ 0 \end{bmatrix} \right) \times \begin{bmatrix} 4 \\ 5 \\ 1 \end{bmatrix} = \mathbf{0}$	<p>M1</p> <p>Sets x, y or $z = \text{constant}$ (eg 0) and attempts to solve the resultant simultaneous equations or Substitutes parametric values to obtain values for x, y and z</p> <p>A1</p> <p>A correct $\mathbf{u}$, eg $\begin{bmatrix} 0 \\ -1 \\ -2 \end{bmatrix}$ or $\begin{bmatrix} 4/5 \\ 0 \\ -9/5 \end{bmatrix}$</p> <p>A1ft</p> <p>Correct for their $\mathbf{v}$ and their $\mathbf{u}$ Must have $= \mathbf{0}$</p>
--	--	---

	Question 12 Total	11	
--	--------------------------	-----------	--

Q	Answer	Marks	Comments
13	Auxiliary equation $m^2 - 2m - 8 = 0$ $m = -2, 4$ $[y_{CF} =] Ae^{-2x} + Be^{4x}$ Particular Integral $[y_{PI} =] Cxe^{-2x} + D$ $[y'_{PI} =] Ce^{-2x} - 2Cxe^{-2x}$ $[y''_{PI} =] -4Ce^{-2x} + 4Cxe^{-2x}$ $(-4Ce^{-2x} + 4Cxe^{-2x})$ $-2(Ce^{-2x} - 2Cxe^{-2x})$ $-8(Cxe^{-2x} + D)$ $= 12e^{-2x} + 8$ e^{-2x} $-6C = 12$ $C = -2$ constant $-8D = 8$ $D = -1$ General Solution $[f(x) =] Ae^{-2x} + Be^{4x} - 2xe^{-2x} - 1$ As $x \rightarrow \infty$ $\lim_{x \rightarrow \infty} (y) = -1$ $= A \lim_{x \rightarrow \infty} (e^{-2x}) + B \lim_{x \rightarrow \infty} (e^{4x}) - 2 \lim_{x \rightarrow \infty} (xe^{-2x}) - 1$ $= B \lim_{x \rightarrow \infty} (e^{4x}) - 1$ $0 = B \lim_{x \rightarrow \infty} (e^{4x})$ $B = 0$ As $\lim_{x \rightarrow \infty} (xe^{-2x}) = 0$ $x = 0$ $y = 3 = A + B - 1 = A - 1$ $A = 4$ $y = 4e^{-2x} - 2xe^{-2x} - 1$	M1 A1 B1 M1 A1 B1 B1 M1 E1 A1 A1	Solves correct auxiliary equation to obtain $m = \pm 2, \pm 4$ Correct CF Correct particular integral. Substitutes their first and second derivatives of y_{PI} and into DE and attempts to compare their coefficients Correct coefficient of e^{-2x} Correct constant term Correct General Solution with exactly two arbitrary constants Uses limits to obtain a value for B Demonstrates $\lim_{x \rightarrow \infty} (xe^{-2x}) = 0$ Correct values for A and B Correct ISW M1 A1 B1 M1 A1 B1 B1 M1 E0 A1 A1 is possible

	Question 13 Total	11	
--	-------------------	----	--

Q	Answer	Marks	Comments
14(a)(i)	$\frac{1}{5} \begin{bmatrix} a & 0 & -2 \\ 0 & 5 & 0 \\ -2 & 0 & 11 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$ $\frac{1}{5} \begin{bmatrix} a-4 \\ 0 \\ 20 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 4 \end{bmatrix}$ $a = 14$	<p>M1</p> <p>A1</p>	<p>Uses $M\mathbf{v}_1 = \lambda_1\mathbf{v}_1$ Or Characteristic equation</p> <p>Correct value of a</p>
		2	

Q	Answer	Marks	Comments
14(a)(ii)	$\frac{1}{5} \begin{bmatrix} 14 & 0 & -2 \\ 0 & 5 & 0 \\ -2 & 0 & 11 \end{bmatrix} \begin{bmatrix} -2 \\ 0 \\ b \end{bmatrix} = \lambda_2 \begin{bmatrix} -2 \\ 0 \\ b \end{bmatrix}$ $\frac{1}{5} \begin{bmatrix} -28-2b \\ 0 \\ 4+11b \end{bmatrix} = \lambda_2 \begin{bmatrix} -2 \\ 0 \\ b \end{bmatrix}$ $\begin{array}{rcl} -28 & -2b & = -10\lambda_2 \\ 4 & +11b & = 5b\lambda_2 \end{array}$ $\frac{-28-2b}{-10} = \frac{4+11b}{5b}$ $10b^2 + 30b - 40 [= 0]$ $b = 1 \text{ [and } b = -4 \text{ (reject as } b > 0)]$ $\lambda_2 = 3$	<p>M1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>Obtains simultaneous equations in b and λ_2</p> <p>Obtains a three-term quadratic expression in b or λ_2 $25\lambda_2^2 - 125\lambda_2 + 150 [= 0]$</p>
		4	

Q	Answer	Marks	Comments
14(a)(iii)	$\frac{1}{5} \begin{bmatrix} 14 & 0 & -2 \\ 0 & 5 & 0 \\ -2 & 0 & 11 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \lambda_3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ $\lambda_3 = 1$	B1	Obtains $\lambda_3 = 1$
		1	

Q	Answer	Marks	Comments
14(b)(i)	$\begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix}$ <p>= 0 so $\mathbf{v}_1$ and $\mathbf{v}_2$ are perpendicular</p> $\begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \times \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix}$ $= \begin{bmatrix} 0 \\ -5 \\ 0 \end{bmatrix} = -5 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ <p>so $\mathbf{v}_1$, $\mathbf{v}_2$ and $\mathbf{v}_3$ are all perpendicular to each other</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Calculates scalar product of two eigenvectors</p> <p>Obtains 0 and interprets this as perpendicular</p> <p>Calculates the vector product of the same two eigenvectors or calculates the scalar products of the other two pairs of eigenvectors</p> <p>Shows that the vector product is a multiple of the third eigenvector and states that the three eigenvectors are all perpendicular to each other or</p> <p>Shows that all three scalar products are zero and states that the three eigenvectors are all perpendicular to each other</p>
		4	

Q	Answer	Marks	Comments
14(b)(ii)	$x = z = 0$	B1	Obtains correct Cartesian equation from $\mathbf{v}_3$
		1	

	Question 14 Total	12	
--	-------------------	----	--

Q	Answer	Marks	Comments
15(a)	$7r^2 + 6(r \sin \theta)^2 + 6\sqrt{3}r \sin \theta r \cos \theta = 64$ $7(x^2 + y^2) + 6y^2 + 6\sqrt{3}yx = 64$ $7x^2 + 6\sqrt{3}xy + 13y^2 = 64$	M1 M1 A1	Correct use of $x = r \cos \theta$ and $y = r \sin \theta$ Correct use of $r^2 = x^2 + y^2$ ACF Condone $64 = 7x^2 + 6\sqrt{3}xy + 13y^2$ NMS = 0/3
		3	

Q	Answer	Marks	Comments
15(b)	$\tan \theta = 1/\sqrt{3} \text{ so } \theta = \pi/6$ $\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos(2\pi/6) & \sin(2\pi/6) \\ \sin(2\pi/6) & -\cos(2\pi/6) \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$ $x = \frac{x' + \sqrt{3} y'}{2}$ $y = \frac{\sqrt{3} x' - y'}{2}$ $64 = 7 \left(\frac{x' + \sqrt{3} y'}{2} \right)^2$ $+ 6\sqrt{3} \left(\frac{x' + \sqrt{3} y'}{2} \right) \left(\frac{\sqrt{3} x' - y'}{2} \right)$ $+ 13 \left(\frac{\sqrt{3} x' - y'}{2} \right)^2$ So equation of curve E is $64 = \frac{1}{4} (7x^2 + 14\sqrt{3} xy + 21y^2) \quad (\text{I})$ $+ \frac{1}{4} (18x^2 + 12\sqrt{3} xy - 18y^2) \quad (\text{II})$ $+ \frac{1}{4} (39x^2 - 26\sqrt{3} xy + 13y^2) \quad (\text{III})$ $64 = 16x^2 + 4y^2$ $16 = 4x^2 + y^2$	<p>B1</p> <p>M1</p> <p>m1</p> <p>A1ft</p> <p>A1</p>	<p>Obtains $x = \frac{x' + \sqrt{3} y'}{2}$ and $y = \frac{\sqrt{3} x' - y'}{2}$ oe Do not condone lack of ‘</p> <p>Uses $x = \frac{\pm x' \pm \sqrt{3} y'}{2}$ and $y = \frac{\pm \sqrt{3} x' \pm y'}{2}$ Condone lack of ‘</p> <p>Correctly expands to obtain at least one of their terms (I), (II) or (III) correct Condone presence of ‘</p> <p>Obtains their correct equation of curve E as the unsimplified sum of x^2, xy and y^2 terms ie their (I), (II) and (III) correct Condone presence of ‘</p> <p>ACF eg $1 = \frac{x^2}{4} + \frac{y^2}{16}$ Condone presence of ‘ NMS = 0/5 B0 M1 m1 A1 A1 is possible</p>
		5	

	Question 15 Total	8	
--	-------------------	---	--